

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
23 August 2001 (23.08.2001)

PCT

(10) International Publication Number  
**WO 01/60805 A1**

(51) International Patent Classification<sup>7</sup>: C07D 239/56,  
A01I 12, 239/70, 239/26, 213/06, 213/54, C07C 15/14,  
A61K 31/505, A61P 3/06

(21) International Application Number: PCT/EP01/01515

(22) International Filing Date: 13 February 2001 (13.02.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
0003636.8 16 February 2000 (16.02.2000) GB  
0101437.2 19 January 2001 (19.01.2001) GB

(71) Applicant (for all designated States except US):  
SMITHKLINE BEECHAM P.L.C. [GB/GB]; New  
Horizons Court, Brentford, Middlesex TW8 9EP (GB).

(72) Inventors; and

(75) Inventors/Applicants (for US only): HICKEY, Deirdre,  
Mary, Bernadette [IE/GB]; GlaxoSmithKline, New  
Frontiers Science Park South, Third Avenue, Harlow  
Essex, CM19 5AW (GB). IFE, Robert, John [GB/GB];  
GlaxoSmithKline, New Frontiers Science Park South,  
Third Avenue, Harlow Essex, CM19 5AW (GB). LEACH,  
Colin, Andrew [GB/GB]; GlaxoSmithKline, New Fron-  
tiers Science Park South, Third Avenue, Harlow Essex,  
CM19 5AW (GB). PINTO, Ivan, Leo [GB/GB]; Glax-  
oSmithKline, New Frontiers Science Park South, Third  
Avenue, Harlow Essex, CM19 5AW (GB). SMITH,

Stephen, Allan [GB/GB]; GlaxoSmithKline, New Fron-  
tiers Science Park South, Third Avenue, Harlow Essex,  
CM19 5AW (GB). STANWAY, Steven, James [GB/GB];  
GlaxoSmithKline, New Frontiers Science Park South,  
Third Avenue, Harlow Essex, CM19 5AW (GB).

(74) Agent: CONNELL, Anthony, Christopher; Corporate  
Intellectual Property, SmithKline Beecham, Two New  
Horizons Court, Brentford, Middlesex TW8 9EP (GB).

(81) Designated States (national): AE, AG, AL, AM, AT, AU,  
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ,  
DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR,  
HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR,  
LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ,  
NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM,  
TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.


(84) Designated States (regional): ARIPO patent (GH, GM,  
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian  
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European  
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,  
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,  
CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

**Published:**

- with international search report
- before the expiration of the time limit for amending the  
claims and to be republished in the event of receipt of  
amendments

For two-letter codes and other abbreviations, refer to the "Guid-  
ance Notes on Codes and Abbreviations" appearing at the begin-  
ning of each regular issue of the PCT Gazette.

(54) Title: PYRIMIDINE-4-ONE DERIVATIVES AS LDL-PLA<sub>2</sub> INHIBITORS



(I)

(57) Abstract: Pyrimidone compounds of formula (I) are inhibitors of  
the enzyme Lp-PLA<sub>2</sub> and are of use in treating atherosclerosis.

WO 01/60805 A1

## Novel Compounds

The present invention relates to certain novel pyrimidinone compounds, processes for their preparation, intermediates useful in their preparation, pharmaceutical compositions containing them and their use in therapy, in particular in the treatment of atherosclerosis.

WO 95/00649 (SmithKline Beecham plc) describe the phospholipase A2 enzyme Lipoprotein Associated Phospholipase A<sub>2</sub> (Lp-PLA<sub>2</sub>), the sequence, isolation and purification thereof, isolated nucleic acids encoding the enzyme, and recombinant host cells transformed with DNA encoding the enzyme. Suggested therapeutic uses for inhibitors of the enzyme included atherosclerosis, diabetes, rheumatoid arthritis, stroke, myocardial infarction, reperfusion injury and acute and chronic inflammation. A subsequent publication from the same group further describes this enzyme (Tew D *et al*, *Arterioscler Thromb Vas Biol* 1996;16:591-9) wherein it is referred to as LDL-PLA<sub>2</sub>. A later patent application (WO 95/09921, Icos Corporation) and a related publication in *Nature* (Tjoelker *et al*, vol 374, 6 April 1995, 549) describe the enzyme PAF-AH which has essentially the same sequence as Lp-PLA<sub>2</sub> and suggest that it may have potential as a therapeutic protein for regulating pathological inflammatory events.

It has been shown that Lp-PLA<sub>2</sub> is responsible for the conversion of phosphatidylcholine to lysophosphatidylcholine, during the conversion of low density lipoprotein (LDL) to its oxidised form. The enzyme is known to hydrolyse the sn-2 ester of the oxidised phosphatidylcholine to give lysophosphatidylcholine and an oxidatively modified fatty acid. Both products of Lp-PLA<sub>2</sub> action are biologically active with lysophosphatidylcholine, a component of oxidised LDL, known to be a potent chemoattractant for circulating monocytes. As such, lysophosphatidylcholine is thought play a significant role in atherosclerosis by being responsible for the accumulation of cells loaded with cholesterol ester in the arteries. Inhibition of the Lp-PLA<sub>2</sub> enzyme would therefore be expected to stop the build up of these macrophage enriched lesions (by inhibition of the formation of lysophosphatidylcholine and oxidised free fatty acids) and so be useful in the treatment of atherosclerosis.

A recently published study (WOSCOPS – Packard *et al*, *N. Engl. J. Med.* 343 (2000) 1148-1155) has shown that the level of the enzyme Lp-PLA<sub>2</sub> is an independent risk factor in coronary artery disease.

The increased lysophosphatidylcholine content of oxidatively modified LDL is also thought to be responsible for the endothelial dysfunction observed in patients with atherosclerosis. Inhibitors of Lp-PLA<sub>2</sub> could therefore prove beneficial in the treatment of this phenomenon. An Lp-PLA<sub>2</sub> inhibitor could also find utility in other disease states that exhibit endothelial dysfunction including diabetes, hypertension, angina pectoris and after ischaemia and reperfusion.

In addition, Lp-PLA<sub>2</sub> inhibitors may also have a general application in any disorder that involves activated monocytes, macrophages or lymphocytes, as all of these cell types express Lp-PLA<sub>2</sub>. Examples of such disorders include psoriasis.


Furthermore, Lp-PLA<sub>2</sub> inhibitors may also have a general application in any disorder that involves lipid oxidation in conjunction with Lp-PLA<sub>2</sub> activity to produce the two injurious products, lysophosphatidylcholine and oxidatively modified fatty acids. Such conditions include the

aforementioned conditions atherosclerosis, diabetes, rheumatoid arthritis, stroke, myocardial infarction, reperfusion injury and acute and chronic inflammation.

Patent applications WO 96/13484, WO96/19451, WO 97/02242, WO97/217675, WO97/217676, WO 97/41098, and WO97/41099 (SmithKline Beecham plc) disclose *inter alia* various series of 4-thionyl/sulfinyl/sulfonyl azetidinone compounds which are inhibitors of the enzyme Lp-PLA<sub>2</sub>. These are irreversible, acylating inhibitors (Tew *et al*, Biochemistry, 37, 10087, 1998).

A further class of compounds has now been identified which are non-acylating inhibitors of the enzyme Lp-PLA<sub>2</sub>. Thus, WO 99/24420 (SmithKline Beecham plc) discloses a class of pyrimidone compounds. International patent applications WO 00/10980, WO 00/66566, WO 00/66567 and WO 00/68208 (SmithKline Beecham plc, published after the priority date of the present application) disclose other classes of pyrimidone compounds. We have now found a further class of pyrimidone compounds which are distinguished by the substitution pattern at the 5 and 6 position of the pyrimidone ring and which have good activity as inhibitors of the enzyme Lp-PLA<sub>2</sub>.

Accordingly, the present invention provides a compound of formula (I):



(I)

in which:

R<sup>a</sup> is hydrogen, halogen, C<sub>(1-3)</sub>alkyl, C<sub>(1-3)</sub>alkoxy, hydroxyC<sub>(1-3)</sub>alkyl, C<sub>(1-3)</sub>alkylthio, C<sub>(1-3)</sub>alkylsulphinyl, aminoC<sub>(1-3)</sub>alkyl, mono- or di-C<sub>(1-3)</sub>alkylaminoC<sub>(1-3)</sub>alkyl, C<sub>(1-3)</sub>alkylcarbonylaminoC<sub>(1-3)</sub>alkyl, C<sub>(1-3)</sub>alkoxyC<sub>(1-3)</sub>alkylcarbonylaminoC<sub>(1-3)</sub>alkyl, C<sub>(1-3)</sub>alkylsulphonylaminoC<sub>(1-3)</sub>alkyl, C<sub>(1-3)</sub>alkylcarboxy, or C<sub>(1-3)</sub>alkylcarboxyC<sub>(1-3)</sub>alkyl;

R<sup>b</sup> is hydrogen, halogen, C<sub>(1-3)</sub>alkyl, or hydroxyC<sub>(1-3)</sub>alkyl, with the proviso that R<sup>a</sup> and R<sup>b</sup> are not simultaneously each hydrogen; or

R<sup>a</sup> and R<sup>b</sup> together are (CH<sub>2</sub>)<sub>n</sub> where n is 3 or 4, to form, with the pyrimidine ring carbon atoms to which they are attached a fused 5- or 6-membered carbocyclic ring; or

R<sup>a</sup> and R<sup>b</sup> together with the pyrimidine ring carbon atoms to which they are attached form a fused benzo or heteroaryl ring optionally substituted by 1, 2, 3 or 4 substituents which may be the same or different selected from halogen, C<sub>(1-4)</sub>alkyl, cyano, C<sub>(1-4)</sub>alkoxy or C<sub>(1-4)</sub>alkylthio, or mono to perfluoro-C<sub>(1-4)</sub>alkyl);

R<sup>c</sup> is hydrogen or C<sub>(1-3)</sub>alkyl;

R<sup>2</sup> is an aryl or heteroaryl group, optionally substituted by 1, 2, 3 or 4 substituents which may be the same or different selected from C<sub>(1-18)</sub>alkyl (preferably C<sub>(1-6)</sub>alkyl), C<sub>(1-18)</sub>alkoxy (preferably C<sub>(1-6)</sub>alkoxy), C<sub>(1-18)</sub>alkylthio (preferably C<sub>(1-6)</sub>alkylthio), arylC<sub>(1-18)</sub>alkoxy (preferably arylC<sub>(1-6)</sub>alkoxy), hydroxy, halogen, CN, COR<sup>6</sup>, carboxy, COOR<sup>6</sup>, NR<sup>6</sup>COR<sup>7</sup>, CONR<sup>8</sup>R<sup>9</sup>,

SO<sub>2</sub>NR<sup>8</sup>R<sup>9</sup>, NR<sup>6</sup>SO<sub>2</sub>R<sup>7</sup>, NR<sup>8</sup>R<sup>9</sup>, mono to perfluoro-C<sub>(1-4)</sub>alkyl, mono to perfluoro-C<sub>(1-4)</sub>alkoxyaryl, and arylC<sub>(1-4)</sub>alkyl;

R<sup>3</sup> is hydrogen, C<sub>(1-6)</sub>alkyl which may be unsubstituted or substituted by 1, 2 or 3 substituents selected from hydroxy, halogen, OR<sup>6</sup>, COR<sup>6</sup>, carboxy, COOR<sup>6</sup>, CONR<sup>8</sup>R<sup>9</sup>, NR<sup>8</sup>R<sup>9</sup>, NR<sup>8</sup>COR<sup>9</sup>, mono- or di-(hydroxyC<sub>(1-6)</sub>alkyl)amino and N-hydroxyC<sub>(1-6)</sub>alkyl-N-C<sub>(1-6)</sub>alkylamino, for instance, 1-piperidinoethyl; or

R<sup>3</sup> is Het-C<sub>(0-4)</sub>alkyl in which Het is a 5- to 7- membered heterocyclyl ring comprising N and optionally O or S, bonded through a carbon ring atom and in which N may be substituted by COR<sup>6</sup>, COOR<sup>6</sup>, CONR<sup>8</sup>R<sup>9</sup>, or C<sub>(1-6)</sub>alkyl optionally substituted by 1, 2 or 3 substituents selected from hydroxy, halogen, OR<sup>6</sup>, COR<sup>6</sup>, carboxy, COOR<sup>6</sup>, CONR<sup>8</sup>R<sup>9</sup> or NR<sup>8</sup>R<sup>9</sup>, for instance, piperidin-4-yl, pyrrolidin-3-yl;

R<sup>4</sup> is an aryl or a heteroaryl ring optionally substituted by 1, 2, 3 or 4 substituents which may be the same or different selected from C<sub>(1-18)</sub>alkyl (preferably C<sub>(1-6)</sub>alkyl), C<sub>(1-18)</sub>alkoxy (preferably C<sub>(1-6)</sub>alkoxy), C<sub>(1-18)</sub>alkylthio (preferably C<sub>(1-6)</sub>alkylthio), arylC<sub>(1-18)</sub>alkoxy (preferably arylC<sub>(1-6)</sub>alkoxy), hydroxy, halogen, CN, COR<sup>6</sup>, carboxy, COOR<sup>6</sup>, NR<sup>6</sup>COR<sup>7</sup>, CONR<sup>8</sup>R<sup>9</sup>, SO<sub>2</sub>NR<sup>8</sup>R<sup>9</sup>, NR<sup>6</sup>SO<sub>2</sub>R<sup>7</sup>, NR<sup>8</sup>R<sup>9</sup>, mono to perfluoro-C<sub>(1-4)</sub>alkyl and mono to perfluoro-C<sub>(1-4)</sub>alkoxy;

R<sup>5</sup> is an aryl or heteroaryl ring which is further optionally substituted by 1, 2, 3 or 4 substituents which may be the same or different selected from C<sub>(1-18)</sub>alkyl (preferably C<sub>(1-6)</sub>alkyl), C<sub>(1-18)</sub>alkoxy (preferably C<sub>(1-6)</sub>alkoxy), C<sub>(1-18)</sub>alkylthio (preferably C<sub>(1-6)</sub>alkylthio), arylC<sub>(1-18)</sub>alkoxy (preferably arylC<sub>(1-6)</sub>alkoxy), hydroxy, halogen, CN, COR<sup>6</sup>, carboxy, COOR<sup>6</sup>, CONR<sup>8</sup>R<sup>9</sup>, NR<sup>6</sup>COR<sup>7</sup>, SO<sub>2</sub>NR<sup>8</sup>R<sup>9</sup>, NR<sup>6</sup>SO<sub>2</sub>R<sup>7</sup>, NR<sup>8</sup>R<sup>9</sup>, mono to perfluoro-C<sub>(1-4)</sub>alkyl and mono to perfluoro-C<sub>(1-4)</sub>alkoxy;

R<sup>6</sup> and R<sup>7</sup> are independently hydrogen or C<sub>(1-20)</sub>alkyl, for instance C<sub>(1-4)</sub>alkyl (e.g. methyl or ethyl);

R<sup>8</sup> and R<sup>9</sup> which may be the same or different is each selected from hydrogen, C<sub>(1-12)</sub>alkyl (preferably C<sub>(1-6)</sub>alkyl); or

R<sup>8</sup> and R<sup>9</sup> together with the nitrogen to which they are attached form a 5- to 7 membered ring optionally containing one or more further heteroatoms selected from oxygen, nitrogen and sulphur, and optionally substituted by one or two substituents selected from hydroxy, oxo, C<sub>(1-4)</sub>alkyl, C<sub>(1-4)</sub>alkylCO, aryl, e.g. phenyl, or aralkyl, e.g. benzyl, for instance morpholine or piperazine; or

R<sup>8</sup> and R<sup>9</sup> which may be the same or different is each selected from CH<sub>2</sub>R<sup>10</sup>, CHR<sup>11</sup>CO<sub>2</sub>H or a salt thereof in which:

R<sup>10</sup> is COOH or a salt thereof, COOR<sup>12</sup>, CONR<sup>6</sup>R<sup>7</sup>, CN, CH<sub>2</sub>OH or CH<sub>2</sub>OR<sup>6</sup>;

R<sup>11</sup> is an amino acid side chain such as CH<sub>2</sub>OH from serine;

R<sup>12</sup> is C<sub>(1-4)</sub>alkyl or a pharmaceutically acceptable *in vivo* hydrolysable ester group;

n is an integer from 1 to 4, preferably 1 or 3, more preferably 1;

X is O or S;

Y is (CH<sub>2</sub>)<sub>p</sub>(O)<sub>q</sub> in which p is 1, 2 or 3 and q is 0 or p is 2 or 3 and q is 1; and

Z is O or a bond.

Representative examples of R<sup>a</sup> include chloro, bromo, methyl, ethyl, n-propyl, methoxy, hydroxymethyl, hydroxyethyl, methylthio, methylsulphinyl, aminoethyl, dimethylaminomethyl, acetylaminomethyl, 2-(methoxyacetamido)ethyl, mesylaminomethyl, ethylcarboxy, methanesulfonamidoethyl, (methoxyacetamido)ethyl and iso-propylcarboxymethyl.

Representative examples of R<sup>b</sup> include hydrogen, and methyl.

Representative examples of  $R^a$  and  $R^b$  together with the pyrimidine ring carbon atoms to which they are attached forming a fused benzo or heteroaryl ring include benzo (to give a quinazolinyl ring), pyrido and thieno, respectively.

- 5 Preferably  $R^a$  is methyl or ethyl and  $R^b$  is hydrogen or methyl, or  $R^a$  and  $R^b$  together with the pyrimidine ring carbon atoms to which they are attached form a fused 5- or 6-membered carbocyclic ring. More preferably,  $R^a$  and  $R^b$  together with the pyrimidine ring carbon atoms to which they are attached form a fused 5-membered carbocyclic ring.

- 10 Representative examples of  $R^c$  include hydrogen and methyl. Preferably,  $R^c$  is hydrogen.

Preferably, X is S.

Preferably, Y is  $CH_2$ .

- 15 Preferably, Z is a direct bond.

- 20 Representative examples of  $R^2$  when an aryl group include phenyl and naphthyl. Representative examples of  $R^2$  when a heteroaryl group include pyridyl, pyrimidinyl, pyrazolyl, furanyl, thienyl, thiazolyl, quinolyl, benzothiazolyl, pyridazolyl and pyrazinyl.

- 25 Preferably,  $R^2$  is an aryl group, optionally substituted by 1, 2, 3 or 4 substituents which may be the same or different selected from  $C_{(1-6)}$ alkyl,  $C_{(1-6)}$ alkoxy,  $C_{(1-6)}$ alkylthio, hydroxy, halogen, CN, mono to perfluoro- $C_{(1-4)}$ alkyl, mono to perfluoro- $C_{(1-4)}$ alkoxyaryl, and aryl $C_{(1-4)}$ alkyl. More preferably,  $R^2$  is phenyl optionally substituted by halogen, preferably from 1 to three fluorine atoms, most preferably 4-fluoro.

Preferably,  $R^2CH_2X$  is 4-fluorobenzylthio.

- 30 Representative examples of  $R^3$  include hydrogen, methyl, 2-(ethylamino)ethyl, 2-(diethylamino)ethyl, 2-(ethylamino)-2-methylpropyl, 2-(t-butylamino)ethyl, 1-piperidinoethyl, 1-ethyl-piperidin-4-yl.


- 35 Preferably,  $R^3$  is  $C_{(1-3)}$ alkyl substituted by a substituent selected from  $NR^8R^9$ ; or  $R^3$  is Het- $C_{(0-2)}$ alkyl in which Het is a 5- to 7- membered heterocyclcyl ring comprising N and in which N may be substituted by  $C_{(1-6)}$ alkyl. More preferably,  $R^3$  is 2-(diethylamino)ethyl.

Representative examples of  $R^4$  include phenyl, pyridine and pyrimidine. Preferably,  $R^4$  is phenyl.

- 40 Representative examples of  $R^5$  include phenyl or thienyl, optionally substituted by halogen or trifluoromethyl, preferably at the 4-position. Preferably,  $R^5$  is phenyl substituted by trifluoromethyl, preferably at the 4-position.

- 45 Preferably,  $R^4$  and  $R^5$  together form a 4-(phenyl)phenyl, 2-(phenyl)pyrimidinyl or a 2-(phenyl)pyridinyl substituent in which the remote phenyl ring may be optionally substituted by halogen or trifluoromethyl, preferably at the 4-position. More preferably,  $R^4$  and  $R^5$  together form a 4-(4-trifluoromethylphenyl)-phenyl moiety.


It will be appreciated that within the compounds of formula (I) there is a sub-group of compounds which has the formula (IA):



in which:

$R^a$ ,  $R^b$ ,  $R^c$ ,  $n$ ,  $R^2$ ,  $R^3$ ,  $R^4$ ,  $R^5$ , and  $X$  are as hereinbefore defined; and

a further sub-group of compounds which has the formula (IB):



in which:

$R^a$ ,  $R^b$ ,  $R^2$ ,  $R^3$ ,  $R^4$ ,  $R^5$ , and  $X$  are as hereinbefore defined, in particular:

$R^a$  and  $R^b$  together with the pyrimidine ring carbon atoms to which they are attached form a fused 5-membered carbocyclic ring;

$R^2CH_2X$  is 4-fluorobenzylthio;

$R^3$  is  $C_{(1-3)}$ alkyl substituted by  $NR^8R^9$ ; or

$R^3$  is Het- $C_{(0-2)}$ alkyl in which Het is a 5- to 7-membered heterocyclyl ring containing N and in which N may be substituted by  $C_{(1-6)}$ alkyl;

$R^4$  and  $R^5$  form a 4-(4-trifluoromethylphenyl)phenyl moiety;

$R^8$  and  $R^9$  which may be the same or different is each selected from hydrogen, or  $C_{(1-6)}$ alkyl; and

$X$  is S.

Pharmaceutically acceptable *in vivo* hydrolysable ester groups for  $R^{12}$  include those which break down readily in the human body to leave the parent acid or its salt. Pharmaceutically acceptable *in vivo* hydrolysable ester groups are well known in the art and examples of such for use in  $R^{12}$  are described in WO 00/68208 (SmithKline Beecham).

It will be appreciated that when R<sup>C</sup> is C<sub>(1-3)</sub>alkyl, the carbon to which it is attached will be a chiral centre so that diastereoisomers may be formed. In the absence of further chiral centres, these will be enantiomers. The present invention covers all such diastereoisomers and enantiomers, including mixtures thereof.

It will be appreciated that in some instances, compounds of the present invention may include a basic function such as an amino group as a substituent. Such basic functions may be used to form acid addition salts, in particular pharmaceutically acceptable salts. Pharmaceutically acceptable salts include those described by Berge, Bighley, and Monkhouse, *J. Pharm. Sci.*, 1977, 66, 1-19. Such salts may be formed from inorganic and organic acids. Representative examples thereof include maleic, fumaric, benzoic, ascorbic, pantoic, succinic, bismethylenesalicylic, methanesulfonic, ethanedisulfonic, acetic, propionic, tartaric, salicylic, citric, gluconic, aspartic, stearic, palmitic, itaconic, glycolic, p-aminobenzoic, glutamic, taurocholic, benzenesulfonic, p-toluenesulfonic, hydrochloric, hydrobromic, sulfuric, cyclohexylsulfamic, phosphoric and nitric acids.

It will be appreciated that in some instances, compounds of the present invention may include a carboxy group as a substituent. Such carboxy groups may be used to form salts, in particular pharmaceutically acceptable salts. Pharmaceutically acceptable salts include those described by Berge, Bighley, and Monkhouse, *J. Pharm. Sci.*, 1977, 66, 1-19. Preferred salts include alkali metal salts such as the sodium and potassium salts.

When used herein, the term "alkyl" and similar terms such as "alkoxy" includes all straight chain and branched isomers. Representative examples thereof include methyl, ethyl, *n*-propyl, *iso*-propyl, *n*-butyl, *sec*-butyl, *iso*-butyl, *t*-butyl, *n*-pentyl and *n*-hexyl.

When used herein, the term "aryl" refers to, unless otherwise defined, a mono- or bicyclic aromatic ring system containing up to 10 carbon atoms in the ring system, for instance phenyl or naphthyl.

When used herein, the term "heteroaryl" refers to a mono- or bicyclic heteroaromatic ring system comprising up to four, preferably 1 or 2, heteroatoms each selected from oxygen, nitrogen and sulphur. Each ring may have from 4 to 7, preferably 5 or 6, ring atoms. A bicyclic heteroaromatic ring system may include a carbocyclic ring.

When used herein, the terms "halogen" and "halo" include fluorine, chlorine, bromine and iodine and fluoro, chloro, bromo and iodo, respectively.

Since the compounds of the present invention, in particular compounds of formula (I), are intended for use in pharmaceutical compositions, it will be understood that they are each provided in substantially pure form, for example at least 50% pure, more suitably at least 75% pure and preferably at least 95% pure (% are on a wt/wt basis). Impure preparations of the compounds of formula (I) may be used for preparing the more pure forms used in the pharmaceutical compositions. Although the purity of intermediate compounds of the present invention is less critical, it will be readily understood that the substantially pure form is preferred as for the compounds of formula (I). Preferably, whenever possible, the compounds of the present invention are obtained in crystalline form.

When some of the compounds of this invention are allowed to crystallise or are re-crystallised from organic solvents, solvent of crystallisation may be present in the crystalline product. This invention includes within its scope such solvates. Similarly, some of the compounds of this invention may be crystallised or re-crystallised from solvents containing water. In such cases water of hydration may be formed. This invention includes within its scope stoichiometric hydrates as well as compounds containing variable amounts of water that may be produced by processes such as lyophilisation. In addition, different crystallisation conditions may lead to the formation of different polymorphic forms of crystalline products. This invention includes within its scope all polymorphic forms of the compounds of formula (I).

Compounds of the present invention are inhibitors of the enzyme lipoprotein associated phospholipase A<sub>2</sub> (Lp-PLA<sub>2</sub>) and as such are expected to be of use in therapy, in particular in the primary and secondary prevention of acute coronary events, for instance those caused by atherosclerosis, including peripheral vascular atherosclerosis and cerebrovascular atherosclerosis. In a further aspect therefore the present invention provides a compound of formula (I) for use in therapy.

The compounds of formula (I) are inhibitors of lysophosphatidylcholine production by Lp-PLA<sub>2</sub> and may therefore also have a general application in any disorder that involves endothelial dysfunction, for example atherosclerosis, diabetes, hypertension, angina pectoris and reperfusion. In addition, compounds of formula (I) may have a general application in any disorder that involves lipid oxidation in conjunction with enzyme activity, for example, in addition to conditions such as atherosclerosis and diabetes, other conditions such as ischaemia, rheumatoid arthritis, stroke, inflammatory conditions of the brain such as Alzheimer's Disease, myocardial infarction, reperfusion injury, sepsis, and acute and chronic inflammation.

Further applications include any disorder that involves activated monocytes, macrophages or lymphocytes, as all of these cell types express Lp-PLA<sub>2</sub>. Examples of such disorders include psoriasis.

Accordingly, in a further aspect, the present invention provides for a method of treating a disease state associated with activity of the enzyme Lp-PLA<sub>2</sub> which method involves treating a patient in need thereof with a therapeutically effective amount of an inhibitor of the enzyme. The disease state may be associated with the increased involvement of monocytes, macrophages or lymphocytes; with the formation of lysophosphatidylcholine and oxidised free fatty acids; with lipid oxidation in conjunction with Lp-PLA<sub>2</sub> activity; or with endothelial dysfunction.

Compounds of the present invention may also be of use in treating the above mentioned disease states in combination with an anti-hyperlipidaemic, anti-atherosclerotic, anti-diabetic, anti-anginal, anti-inflammatory, or anti-hypertension agent or an agent for lowering Lp(a). Examples of the above include cholesterol synthesis inhibitors such as statins, anti-oxidants such as probucol, insulin sensitisers, calcium channel antagonists, and anti-inflammatory drugs such as NSAIDs. Examples of agents for lowering Lp(a) include the aminophosphonates described in WO 97/02037, WO 98/28310, WO 98/28311 and WO 98/28312 (Symphar SA and SmithKline Beecham).

It is expected that compounds of the present invention may be used in combination with cholesterol lowering agents, for instance co-administered with a statin. The statins are a well known class of cholesterol lowering agents (HMG-CoA reductase inhibitors) and include atorvastatin, simvastatin,



pravastatin, cerivastatin, fluvastatin, lovastatin and ZD 4522 (also referred to as S-4522, Astra Zeneca). The two agents may be administered at substantially the same time or at different times, according to the discretion of the physician.

- 5 A substantial minority (approx 30%) of patients with elevated levels of cholesterol are found to not respond to treatment with a statin. In a further use, a compound of the present invention is administered to a patient who has failed to respond to treatment with a statin.

- 10 A further preferred combination therapy will be the use of a compound of the present invention and an anti-diabetic agent or an insulin sensitiser, as coronary heart disease is a major cause of death for diabetics. Within this class, preferred compounds for use with a compound of the present invention include the PPARgamma activators, for instance GI262570 (Glaxo Wellcome) and the glitazone class of compounds such as rosiglitazone (Avandia, SmithKline Beecham), troglitazone and pioglitazone.

- 15 Preferred indications include primary and secondary prevention of acute coronary events, for instance those caused by atherosclerosis, including peripheral vascular atherosclerosis and cerebrovascular atherosclerosis; adjunctive therapy in prevention of restenosis, and delaying the progression of diabetic / hypertensive renal insufficiency.


- 20 In therapeutic use, the compounds of the present invention are usually administered in a standard pharmaceutical composition. The present invention therefore provides, in a further aspect, a pharmaceutical composition comprising a compound of formula (I) and a pharmaceutically acceptable carrier.

- 25 Suitable pharmaceutical compositions include those which are adapted for oral or parenteral administration or as a suppository. Compounds of formula (I) which are active when given orally can be formulated as liquids, for example syrups, suspensions or emulsions, tablets, capsules and lozenges. A liquid formulation will generally consist of a suspension or solution of the compound or pharmaceutically acceptable salt in a suitable liquid carrier(s) for example, ethanol, glycerine, non-aqueous solvent, for example polyethylene glycol, oils, or water with a suspending agent, preservative, flavouring or colouring agent. A composition in the form of a tablet can be prepared using any suitable pharmaceutical carrier(s) routinely used for preparing solid formulations. Examples of such carriers include magnesium stearate, starch, lactose, sucrose and cellulose. A composition in the form of a capsule can be prepared using routine encapsulation procedures. For example, pellets containing the active ingredient can be prepared using standard carriers and then filled into a hard gelatin capsule; alternatively, a dispersion or suspension can be prepared using any suitable pharmaceutical carrier(s), for example aqueous gums, celluloses, silicates or oils and the dispersion or suspension then filled into a soft gelatin capsule. Typical parenteral compositions consist of a solution or suspension of the compound of formula (I) in a sterile aqueous carrier or parenterally acceptable oil, for example polyethylene glycol, polyvinyl pyrrolidone, lecithin, arachis oil or sesame oil. Alternatively, the solution can be lyophilised and then reconstituted with a suitable solvent just prior to administration. A typical suppository formulation comprises a compound of formula (I) which is active when administered in this way, with a binding and/or lubricating agent such as polymeric glycols, gelatins or cocoa butter or other low melting vegetable or synthetic waxes or fats.

45

Preferably the composition is in unit dose form such as a tablet or capsule. Each dosage unit for oral administration contains preferably from 1 to 500 mg (and for parenteral administration contains preferably from 0.1 to 25 mg) of a compound of the formula (I). The daily dosage regimen for an adult patient may be, for example, an oral dose of between 1 mg and 1000 mg, preferably between 1 mg and 500 mg, or an intravenous, subcutaneous, or intra-muscular dose of between 0.1 mg and 100 mg, preferably between 0.1 mg and 25 mg, of the compound of the formula (I), the compound being administered 1 to 4 times per day. Suitably the compounds will be administered for a period of continuous therapy, for example for a week or more.

- 10 A compound of formula (I) may be prepared by reacting a compound of formula (II):



(II)

- 15 in which X, n, Ra, Rb and R2 are as hereinbefore defined,  
with a compound of formula (III):




(III)

in which R<sup>c</sup>, R<sup>3</sup>, R<sup>4</sup>, R<sup>5</sup>, Y and Z are as hereinbefore defined; under amide forming conditions.

Amide forming conditions are well known in the art, see for instance Comprehensive Organic Synthesis 6, 382-399, and include reacting the acid compound of formula (II) and the amine compound of formula (III) in an inert solvent such as dichloromethane, at ambient temperature, in the presence of a coupling agent. Preferred coupling agents include those developed for use in peptide chemistry, such as 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride ("EDC"), preferably in the presence of an additive such as 1-hydroxybenzotriazole, or O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate ("HATU"), preferably in the presence of di-isopropylethylamine.

Compounds of formula (I) may also be prepared by a number of other processes, for instance:


- (a) reacting a compound of formula (IV):



(IV)

- in which X, Ra, Rb and R2 are as hereinbefore defined,


with a compound of formula (V):



(V)

- 5 in which  $n$ ,  $R^3$ ,  $R^4$ ,  $R^5$ ,  $R^c$ ,  $Y$  and  $Z$  are as hereinbefore defined, and  $L^1$  is a leaving group such as halogen, for instance bromo iodo, or triflate;  
in the presence of a base such as a secondary or tertiary amine, for instance di-isopropylethylamine, in an inert solvent such as dichloromethane;

- 10 (b) when  $X$  is  $S$ , reacting a compound of formula (VI):



(VI)


- 15 in which  $n$ ,  $R^a$ ,  $R^b$ ,  $R^c$ ,  $R^3$ ,  $R^4$ ,  $R^5$ ,  $Y$  and  $Z$  are as hereinbefore defined,  
with a compound of formula (VII):



(VII)

- in which  $R^2$  and  $L^1$  are as hereinbefore defined,  
20 in the presence of a base such as a secondary or tertiary amine, for instance di-isopropylethylamine, in an inert solvent such as dichloromethane; or

(c) when  $X$  is  $O$ , reacting a compound of formula (VIII):



(VIII)

25

in which  $n$ ,  $R^a$ ,  $R^b$ ,  $R^c$ ,  $R^3$ ,  $R^4$ ,  $R^5$ ,  $Y$  and  $Z$  are as hereinbefore defined, and  $L^2$  is a leaving group such as halogen or alkylthio, for instance methylthio, with a compound of formula (IX):



(IX)


in which  $R^2$  is as hereinbefore defined,

in the presence of a base such as 4-dimethylaminopyridine, in an inert solvent such as pyridine.

- 10 It will be appreciated that an initially prepared compound of formula (I) may be converted to another compound of formula (I), by functional group modification, using methods well known to those skilled in the art, for example converting a compound of formula (I) in which  $R^a$  is aminoalkyl to a compound of formula (I) in which  $R^a$  is alkylcarbonylaminoalkyl, by reaction with an acylating agent, such as, for example, acetic anhydride.

15

Compounds of formulae (II), (IV), (VI) and (VIII) for use in the above processes may be prepared by processes illustrated in the following scheme I:



Scheme I

in which:

L<sup>3</sup> is a C(1-6)alkyl group, for instance methyl;

R<sup>15</sup> is a C(1-6)alkyl group, for instance ethyl or t-butyl and

L<sup>1</sup>, L<sup>2</sup>, R<sup>a</sup>, R<sup>b</sup>, R<sup>c</sup>, R<sup>2</sup>, R<sup>3</sup>, R<sup>4</sup>, R<sup>5</sup>, n, X, Y and Z are as hereinbefore defined.

With reference to Scheme I:

Amide forming conditions for step (a) are well known in the art. Preferably, the acid of formula (II) is reacted with the amine of formula (III) in an inert solvent, such as dichloromethane, at ambient temperature and in the presence of a coupling agent such as O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate and di-isopropylethylamine or 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride in the presence of 1-hydroxybenzotriazole.

Alkylation conditions for step (b) include reaction in the presence of a base such as a secondary or tertiary amine, for instance di-isopropylethylamine, in an inert solvent such as dichloromethane.

Conditions for step (c) include hydrolysis, for instance using aqueous sodium hydroxide in a solvent such as dioxan or, when R<sup>15</sup> is t-butyl, dealkylation with an acid such as trifluoroacetic acid in a solvent such as dichloromethane.

Conditions for step (d) include under thioether forming conditions. Advantageously, the reaction is carried out in the presence of a base such as sodium ethoxide or potassium carbonate, preferably in a solvent such as ethanol, dimethyl formamide or acetone, or a secondary or tertiary amine base such as di-isopropylethylamine, in solvent such as dichloromethane.

In step (e), a compound of formula (XVII) is reacted with thiourea, in the presence of sodium ethoxide (preferably generated *in situ* from sodium and ethanol).

In step (f), a compound of formula (XVIII) is reacted with ethyl formate in the presence of a base such as sodium hydride or potassium iso-propoxide.

In step (g), a compound of formula (IV) is reacted with a compound of formula (V) in the presence of a base such as a secondary or tertiary amine, for instance di-isopropylethylamine, in an inert solvent such as dichloromethane

In step (h), a compound of formula (XIII) is reacted with a compound of formula (XIV) in a solvent such as dimethylformamide to form an intermediate thiourea, which is then treated with a base such as sodium methoxide.

In step (i), a compound of formula (XVI) is reacted with a metal thiocyanate, for example potassium thiocyanate, in a solvent such as acetonitrile.

In step (j), a compound of formula (XVII) is reacted with a methylating agent such as dimethyl sulphate in the presence of a base such as potassium carbonate, followed by hydrolysis of the intermediate ester in conventional manner e.g. by basic hydrolysis using sodium hydroxide to give the corresponding

carboxylic acid which may then be converted into the acyl chloride, for instance by treatment with oxalyl chloride.

In step (k), a catalyst such as 4-dimethylaminopyridine, and in a solvent such as pyridine are used.

5

In step (l), a compound of formula (XIII) is reacted with a compound of formula (XV) in a solvent such as dimethylformamide to form an intermediate thiourea, which is then treated with a base such as sodium methoxide.

10 In step (m) a compound of formula (XX) is converted to a compound of formula (XIX), in which  $R^a$  is halogen, by treatment with N-halosuccinimide, for example N-chlorosuccinimide or N-bromosuccinimide, in a solvent such as carbon tetrachloride.


15 Compounds of formula (II) and (IV), in particular wherein  $R^a$  and  $R^b$  together with the pyrimidine ring carbon atoms to which they are attached form a fused 5-membered carbocyclic ring, are novel and form a further aspect of the present invention.

The present invention will now be illustrated by the following examples.

## Examples

The structure and purity of the intermediates and examples was confirmed by <sup>1</sup>H-NMR and (in nearly all cases) mass spectroscopy, even where not explicitly indicated below.


## 5 Intermediate A1 — 4-(4-Chlorophenyl)benzaldehyde



(a) A mixture of 4-formylbenzeneboronic acid (2.50g, 2 equiv), 4-chloriodobenzene (1.98g, 1 equiv), tetrakis(triphenylphosphine)palladium(0) (0.50g, 0.05 equiv), aqueous sodium carbonate (18ml, 2M solution, 2 equiv) and dimethoxyethane (50ml) was stirred at reflux under argon overnight, then cooled and diluted with ethyl acetate. The mixture was filtered as necessary to remove inorganic residues, then the organic layer was washed successively with aqueous citric acid and brine, dried and evaporated. The crude product was purified by chromatography (silica, 5% ethyl acetate in hexane); product fractions were evaporated to a white solid (1.32g, 72%).

(b) A mixture of 4-chlorobenzeneboronic acid (19.4g, 1 equiv), 4-bromobenzaldehyde (22.9g, 1 equiv), palladium(II) acetate (1.4g, 0.05 equiv) aqueous sodium carbonate (30.3 g in 144ml solution, 2 equiv) and dimethoxyethane (500ml) was stirred at reflux under argon for 2.5h, then evaporated to low volume and diluted with dichloromethane. Workup continued as in (a) above to give identical material (25.2g, 94%). <sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ 10.05 (1H, s), 7.96 (2H, d), 7.73 (2H, d), 7.57 (2H, d), 7.46 (2H, d); MS (AP+) found (M+1) = 217, C<sub>13</sub>H<sub>9</sub><sup>35</sup>ClO requires 216.


## 20 Intermediate A2 — N-Methyl-4-(4-chlorophenyl)benzylamine



A mixture of Intermediate A1 (3.5g, 1 equiv), methylamine (32.3ml of a 2M solution in THF, 4 equiv) and anhydrous magnesium sulphate (4.47g, 2 equiv) was stirred at room temperature for 16h, then filtered, the solid washed thoroughly with ethyl acetate, and the combined filtrates evaporated to a white solid

(3.7g). This imine intermediate was suspended in ethanol (100ml), cooled in ice and sodium borohydride (0.61g, 1 equiv) added portionwise. The ice bath was removed, and the mixture stirred for 45min at room temperature then at 50°C for 1h. The solvent was removed in vacuo, water was added to the residue, and the product extracted into dichloromethane. Drying and evaporation of the solvent gave a white solid (3.56g). <sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ 7.51 (4H, d), 7.40 (4H, d), 3.79 (2H, s), 2.48 (3H, s); MS (APCI+) found (M+1) = 232, C<sub>14</sub>H<sub>14</sub><sup>35</sup>ClN requires 231.


## Intermediate A3 — N-(2-Diethylaminoethyl)-4-(4-chlorophenyl)benzylamine



A mixture of Intermediate A1 (55.0g), N,N-diethylethylenediamine (35.6ml), 4A molecular sieve (37g), and dichloromethane (1100ml) was reacted at room temperature under argon for 16h, with occasional agitation. The solid was filtered off and washed with dichloromethane, and the combined filtrates evaporated to a yellow foam (72.4g). This intermediate imine was reduced with sodium borohydride (8.7g) in ethanol (850ml) as described for Intermediate A2, yielding the title compound as a yellow oil


(72.7g).  $^1\text{H-NMR}$  ( $\text{CDCl}_3$ )  $\delta$  1.70 (2H, t), 2.22 (6H, s), 2.33 (2H, t), 2.69 (2H, br. m), 3.83 (2H, s), 7.37-7.43 (4H, m), 7.52-7.56 (4H, m).

**Intermediate A4 — 5-Hydroxymethyl-2-(4-trifluoromethylphenyl)pyridine**




- 5 A solution of Intermediate A20 (4.63g) in dry dichloromethane (100ml) was cooled to  $-78^\circ\text{C}$  under argon, then DIBAL-H (26.7ml, 1.5M solution in toluene) was added dropwise over 20min. Stirring was continued for 40min at  $-78^\circ\text{C}$ , then 2M hydrochloric acid (52ml) was added dropwise over 15min. The solution was allowed to warm slowly to room temperature, then the organic layer was separated, washed with water, dried and evaporated. Chromatography (silica, 1:1 ethyl acetate/hexane) gave the product as
- 10 a white solid (3.03g, 75%).  $^1\text{H-NMR}$  ( $\text{CDCl}_3$ )  $\delta$  1.85 (1H,t), 4.81 (2H,d), 7.75 (2H,m), 7.83 (1H,dd), 8.11 (1H,d), 8.72 (1H,m); MS(APCI+) found (M+1)=254,  $\text{C}_{13}\text{H}_{10}\text{F}_3\text{NO}$  requires 253.

**Intermediate A5 — 5-Formyl-2-(4-trifluoromethylphenyl)pyridine**




- 15 Activated manganese dioxide (3.19g) was added to a solution of Intermediate A4 (0.75g) in dichloromethane (50ml) and stirred at room temperature for 16h. The solids were filtered off and the filtrate evaporated to a pale yellow solid (0.57g).  $^1\text{H-NMR}$  ( $\text{CDCl}_3$ )  $\delta$  7.7 (2H,d), 7.96 (1H,d), 8.21 (2H,d), 8.27 (1H,dd), 9.17 (1H,d), 10.19 (1H,s); MS(APCI+) found (M+1)=252,  $\text{C}_{13}\text{H}_8\text{F}_3\text{NO}$  requires 251.

**Intermediate A6 — Ethyl 2-(4-chlorophenyl)-4-oxopyrimidine-5-carboxylate**




- 20 Sodium ethoxide (11.12 ml, 2 equiv) as a 21% w/v solution in ethanol was added dropwise to a suspension of diethyl ethoxymalonate (3.03 ml, 1 equiv) and 4-chlorobenzamidine hydrochloride (4.23 g, 1 equiv) in ethanol (30 ml), then the mixture was heated to reflux for 4 hours. After cooling, the solvent was removed in vacuo and the residue was triturated with ether. The solid was filtered off, then
- 25 resuspended in water and acidified to pH 2. The product was filtered off, washed with water and dried; yield 2.94 g.  $^1\text{H-NMR}$  ( $d_6$ -DMSO)  $\delta$  1.29 (3H,t), 4.26 (2H,q), 7.65 (2H,m), 8.18 (2H,m), 8.65 (1H,s); MS (APCI-) found (M-1) = 277/279;  $\text{C}_{13}\text{H}_{11}\text{ClN}_2\text{O}_3$  requires 278/280.

**Intermediate A7 — Ethyl 2-(4-chlorophenyl)-4-chloropyrimidine-5-carboxylate**




- 30 Oxalyl chloride (0.31 ml, 2 equiv) was added to Intermediate A6 (0.49 g) in dichloromethane (20 ml) with ice cooling, then the mixture was stirred for 3 hours with warming to room temperature. Evaporation of the volatile components gave the product as a white solid (2.94 g).  $^1\text{H-NMR}$  ( $\text{CDCl}_3$ )  $\delta$  1.44 (3H,t), 4.48 (2H,q), 7.50 (2H,m), 8.45 (2H,m), 9.17 (1H,s); MS (APCI+) found (M+1) = 297;  $\text{C}_{13}\text{H}_{10}\text{Cl}_2\text{N}_2\text{O}_2$  requires 296.




**Intermediate A8 — Ethyl 2-(4-chlorophenyl)pyrimidine-5-carboxylate**


A mixture of Intermediate A7 (6.8 g, 1 equiv), zinc powder (1.79 g, 1.2 equiv), acetic acid (1.57 ml, 1.2 equiv) and THF (100 ml) was stirred at 60°C under argon for 18 hours, then a further portion of acetic acid (1 ml) and zinc (1.0 g) was added, and the reaction allowed to continue for a further 24 hours. The solvent was removed in vacuo, the residue was taken up in a mixture of dichloromethane and methanol, and undissolved zinc powder was removed by filtration. After evaporation of the solvent, the product crystallised from ethanol; yield 2.02 g. <sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ 1.44 (3H,t), 4.46 (2H,q), 7.48 (2H,m), 8.48 (2H,m), 9.30 (2H,s); MS (APCI+) found (M+1) = 263; C<sub>13</sub>H<sub>11</sub>ClN<sub>2</sub>O<sub>2</sub> requires 262.

**Intermediate A9 — 5-Hydroxymethyl-2-(4-trifluoromethylphenyl)pyrimidine**


Intermediate A41 (0.96g) was hydrogenated over 10% palladium on charcoal (96 mg) in a mixture of triethylamine (2 ml) and ethanol (20 ml) for 90 mins at 1 atmosphere pressure. The catalyst was removed by filtration, the solvent was evaporated, and the residue was taken up in ethyl acetate and washed successively with aq. ammonium chloride and aq. sodium bicarbonate. Drying and evaporation gave the title compound (0.77 g). <sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ 4.82 (2H,s), 7.75 (2H,m), 8.57 (2H,m), 8.85 (2H,s); MS (APCI+) found (M+1) = 255; C<sub>12</sub>H<sub>9</sub>F<sub>3</sub>N<sub>2</sub>O requires 254.

**Intermediate A10 — 3-(4-trifluoromethylphenoxy)benzyl alcohol**


A mixture of 4-chlorobenzotrifluoride (27.1 g, 1.5 equiv), 3-hydroxybenzyl alcohol (12.4 g, 1 equiv), copper (I) chloride (0.2 g, 0.02 equiv), potassium carbonate (8.3 g, 0.6 equiv), 8-quinolinol (0.29 g, 0.02 equiv) and 1,3-dimethyl-2-imidazolidinone (50 mL) was stirred at 150°C under argon for 3 days. After cooling, the residue was poured into water and extracted with ethyl acetate. Drying and evaporation, followed by chromatography (silica, dichloromethane) gave the title compound as a pale liquid (11.3 g). <sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ 1.88 (1H,t), 4.69 (2H,d), 6.97 (1H,m), 7.04 (3H,m), 7.17 (1H,m), 7.36 (1H,m), 7.57 (2H,m); MS (APCI-) found (M-1) = 267; C<sub>14</sub>H<sub>11</sub>F<sub>3</sub>O<sub>2</sub> requires 268.

**Intermediate A11 — 4-(4-trifluoromethylphenoxy)benzaldehyde**


A mixture of 4-(trifluoromethyl)phenol (4.86 g, 1 equiv), 4-fluorobenzaldehyde (3.22 mL, 1 equiv), potassium carbonate (4.15 g, 1 equiv) and dimethylformamide (60 mL) was stirred at 150°C under argon for 3 hours, then poured into ice/water. The precipitate was filtered off, washed with water, then extracted with hot ethanol. Undissolved solid was removed by filtration, and the filtrate evaporated and purified by chromatography on silica. <sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ 7.14 (4H,m), 7.66 (2H,m), 7.90 (2H,m), 9.97 (1H,s); MS (APCI+) found (M+1) = 267; C<sub>14</sub>H<sub>9</sub>F<sub>3</sub>O<sub>2</sub> requires 266.

**Intermediate A12 — *tert*-Butyl (2-hydroxyethyl)ethylcarbamate**


- 5 Di-*tert*-butyl dicarbonate (15.5g, 1 equiv) was added over a period of 1 hour to a solution of 2-(ethylamino)ethanol (7.5g, 1 equiv) in dichloromethane (30ml) at 0°C. After stirring at room temperature for 16 hours, the solvent was evaporated and the residue distilled (115°C, 0.6mmHg) to afford the title compound as a colourless oil (13.42g). <sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ 1.11 (3H,t), 1.47 (9H,s), 3.27 (2H,q), 3.38 (2H,t), 3.75 (2H,t).

**Intermediate A13 — *tert*-Butyl [2-(phthalimidyl)ethyl]ethylcarbamate**


- 10 Diethylazodicarbonate (12.35g, 1 equiv) was added dropwise to a mixture of intermediate A12 (13.42g, 1 equiv), phthalimide (10.43g, 1 equiv) and triphenylphosphine (18.6g, 1 equiv) in THF (200ml) and the mixture stirred at room temperature for 16 hours. The solvent was evaporated and diethyl ether added. The solution was cooled to 0°C and the insoluble products removed by filtration. The solvent was evaporated and the residue applied to a column (silica, 9:1 Hexane/ethyl acetate) to afford the title compound as a colourless oil (17g). <sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ 1.13 (3H,m), 1.29 (9H,s), 3.26 (2H,m), 3.48 (2H,m), 3.84 (2H,t), 7.71 (2H,m), 7.85 (2H,m).

**Intermediate A14 — *tert*-Butyl (2-Aminoethyl)ethylcarbamate**


- 20 Hydrazine monohydrate (5.2ml, 2 equiv) was added to a solution of intermediate A13 (17g, 1 equiv) in ethanol (300ml) and the reaction stirred for 16 hours at room temperature. The resultant solid was filtered off and the solvent evaporated. The residue was partitioned between diethyl ether and sodium hydroxide (1M, 150ml) and the organic phase dried (K<sub>2</sub>CO<sub>3</sub>) and the solvent removed to afford the title compound as a yellow oil (9.05g). <sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ 1.10 (3H,t), 1.45 (9H,s), 2.65 (2H,q), 2.73 (2H,t), 3.23 (2H,m).

**Intermediate A15 — 3-(4-Trifluoromethyl-biphenyl-4-yl)propan-1-ol**


- 30 Borane in tetrahydrofuran (1.0M, 44.5ml, 2.5 equiv) was added dropwise to a solution of intermediate A23 (5.23g, 1 equiv) in tetrahydrofuran (65ml) at 0°C. The solution was allowed to warm to room temperature and stirring continued for 16 hours. The reaction was quenched by the addition of water and the mixture extracted with ethyl acetate. The organic phase was washed with aq. sodium bicarbonate, dried (MgSO<sub>4</sub>) and the solvent evaporated to afford a residue which was applied to a column (silica, dichloromethane) to afford the title compound as a colourless solid (4.31g). <sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ 1.76 (2H,m), 2.67 (2H,t), 3.45 (2H,m), 7.32 (2H,d), 7.64 (2H,d), 7.78 (2H,d), 7.86 (2H,d).

**Intermediate A16 — 3-(4-Trifluoromethyl-biphenyl-4-yl)propionaldehyde**


Dimethylsulphoxide (2.36ml, 2.4 equiv) was added dropwise to a solution of oxalyl chloride (1.46ml, 1.1equiv) in dichloromethane (34ml) at  $-55^{\circ}\text{C}$  and the solution stirred for 2 minutes. A solution of intermediate A15 (4.28g, 1 equiv) in dichloromethane (40ml) was added slowly to the solution at  $-55^{\circ}\text{C}$  and the solution stirred for a further 10 minutes prior to the addition of triethylamine (9.7ml, 5 equiv). After stirring for a further 5 minutes the reaction was allowed to warm to room temperature and then diluted with water. The organic phase was separated, dried ( $\text{MgSO}_4$ ) and the solvent removed to afford the title compound (3.48g).  $^1\text{H-NMR}$  ( $\text{CDCl}_3$ )  $\delta$  2.83 (2H,m), 3.02 (2H,t), 7.29 (2H,d), 7.51 (2H,d), 7.67 (4H,s), 9.85 (1H,s). MS (APCI+) found ( $\text{M}+1$ ) = 279;  $\text{C}_{16}\text{H}_{13}\text{F}_3\text{O}$  requires 278.

**Intermediate A17 — C-(4'-Trifluoromethyl-biphenyl-4-yl)methylamine**


A solution of intermediate A130 (31g, 1 equiv) in tetrahydrofuran (300ml) was added dropwise to a solution of lithium aluminium hydride (1.0M in tetrahydrofuran, 188ml, 1.5 equiv) at room temperature with stirring. The reaction was stirred for 8 hours, after which time aq. ammonium chloride (200ml) and then water (200ml) was added. The resultant mixture was filtered through celite and then extracted with dichloromethane. The organic phase was dried ( $\text{MgSO}_4$ ) and solvent removed to afford the title compound (26.7g).  $^1\text{H-NMR}$  (DMSO)  $\delta$  3.89 (2H,s), 7.52 (2H,d), 7.73 (2H,d), 7.82 (2H,d), 7.98 (2H,d).

**Intermediate A18 — N-(1-Ethyl-piperidin-4-yl)-(4'-trifluoromethylphenyl)benzylamine**


A solution of intermediate A17 (9.3g, 1 equiv) and 1-ethyl-4-piperidone (5.0ml, 1.05 equiv) in 1,2-dichloroethane (135ml) was treated with sodium triacetoxymethylborohydride (11g, 1.4 equiv) and acetic acid (2.23g, 1.05 equiv) at room temperature and the mixture was stirred for 24 hours. The reaction was quenched by the addition of sodium hydroxide (2M, 125ml) and extracted with diethyl ether. The organic phase was dried ( $\text{MgSO}_4$ ) and solvent evaporated to afford a residue, which was triturated with hexane to afford the title compound as a off white solid (8.2g).  $^1\text{H-NMR}$  ( $\text{CDCl}_3$ )  $\delta$  1.06 (3H,t), 1.48 (3H,m), 2.01 (4H,m), 2.38 (2H,q), 2.55 (1H, m), 2.92 (2H,m), 3.88 (2H,s), 7.43 (2H,d), 7.59 (2H,d), 7.68 (4H,s).

**Intermediate A120 — *tert*-Butyl (2-Amino-2-methylpropyl)-carbamate**


Di-*tert*-butyl dicarbonate (6.58g, 1 equiv) in tetrahydrofuran (100ml) was added dropwise to a solution of 1,2-diamino-2-methylpropane (8.86g, 3.3 equiv) in tetrahydrofuran (100ml) at  $0^{\circ}\text{C}$ . The solution was then stirred at room temperature for 16 hours. The solvent was evaporated and the residue partitioned between aq. sodium chloride and ethyl acetate. The organic phase was dried ( $\text{K}_2\text{CO}_3$ ) and solvent evaporated to afford the title compound as a colourless solid (5.45g).  $^1\text{H-NMR}$  ( $\text{CDCl}_3$ )  $\delta$  1.09 (6H,s), 1.45 (9H,s), 3.00 (2H,d). MS (APCI+) found ( $\text{M}+1$ ) = 189;  $\text{C}_9\text{H}_{20}\text{N}_2\text{O}_2$  requires 188.

**Intermediate A121 — *tert*-Butyl (2-Ethylamino-2-methylpropyl)-carbamate**


Intermediate A120 (5.45g, 1 equiv), iodoethane (2.32ml, 1 equiv) and potassium carbonate (4g, 1 equiv) in dimethylformamide (80ml) were stirred at room temperature for 16 hours. Solvent was evaporated and the residue partitioned between dichloromethane and water. The organic layer was dried ( $K_2CO_3$ ), solvent evaporated and the residue applied to a column (silica, 10:1 dichloromethane/methanol) to afford the title compound as a light brown oil (3.89g).  $^1H$ -NMR ( $CDCl_3$ )  $\delta$  1.05 (6H,s), 1.08 (3H,t), 1.45 (9H,s), 2.54 (2H,q), 3.03 (2H,m). MS (APCI+) found  $(M+1) = 217$ ;  $C_{11}H_{24}N_2O_2$  requires 216.

**Intermediate A122 —  $N^2$ -Ethyl-2-methylpropane-1,2-diamine dihydrochloride**

Hydrogen chloride (4M in dioxan, 70ml) was added to a solution of intermediate A121 (3.89g) in dioxan (100ml) and the resulting suspension stirred at room temperature for 16 hours. Solvent was evaporated and the residue suspended in diethyl ether, the resulting solid was filtered off and collected to afford the title compound as a colourless solid (2.99g).  $^1H$ -NMR ( $d_6$ -DMSO)  $\delta$  1.26 (3H,t), 1.39 (6H,s), 2.97 (2H,q), 3.19 (2H,s). MS (APCI+) found  $(M+1) = 117$ ;  $C_6H_{16}N_2$  requires 116.

**Intermediate A123 — 2-(2-*tert*-Butylaminoethyl)phthalimide**

A mixture of 2-bromoethyl phthalimide (20g, 2 equiv), *tert*-butylamine (41ml, 1 equiv) and potassium carbonate (10.86g, 2 equiv) in dimethylformamide (200ml) was heated to 50°C for 48 hours. Solvent was evaporated and the residue partitioned between dichloromethane and water. The organic phase was dried ( $K_2CO_3$ ) and solvent removed to afford the title compound as an orange solid (18.93g).  $^1H$ -NMR ( $CDCl_3$ )  $\delta$  1.05 (9H,s), 2.85 (2H,t), 3.77 (2H,t), 7.72 (2H,m), 7.85 (2H,m).

**Intermediate A124 — *N*-*tert*-Butylethane-1,2-diamine**

A mixture of intermediate A123 (4g, 1 equiv) and hydrazine hydrate (1.58ml, 2 equiv) in methylated spirit (100ml) was heated to reflux for 16 hours. Solid filtered off and solution used directly in the next step.

The following intermediates were made by the method of Intermediate A1:

No.	Precursors	Name
A20	methyl 6-chloronicotinate, 4-trifluoromethylbenzeneboronic acid	Methyl 6-(4-trifluoromethylphenyl)nicotinate
A21	4-bromobenzaldehyde, 4-trifluoromethylbenzeneboronic acid	4-(4-Trifluoromethylphenyl)benzaldehyde
A22	4-bromoacetophenone, 4-chlorobenzeneboronic acid	4-acetyl-4'-chlorobiphenyl



<b>A23</b>	4-(trifluoromethyl)bromobenzene 4-(2-carboxyethyl)phenylboronic acid	3-(4-Trifluoromethyl-biphenyl-4-yl)propionic acid
<b>A24</b>	2-(4-bromophenoxy)ethanol 4-trifluoromethylbenzeneboronic acid	2-(4-trifluoromethyl-biphenyloxy)ethanol
<b>A130</b>	4-bromobenzonitrile 4-trifluoromethylbenzeneboronic acid	4'-trifluoromethyl-biphenyl-4-carbonitrile

The following intermediates were made by the method of Intermediate A2:

No.	Precursor	Structure	Name
<b>A25</b>	Int. A21		N-Methyl-4-(4-trifluoromethylphenyl)benzylamine
<b>A26</b>	Int. A5		N-methyl-2-(4-trifluoromethylphenyl)pyrid-5-yl-methylamine

The following intermediates were made by the method of Intermediate A3:

No.	Precursor	Structure	Name
<b>A30</b>	Int. A21		N-(2-(diethylamino)ethyl)-4-(4-trifluoromethylphenyl)benzylamine
<b>A31</b>	Int. A5		N-(2-(diethylamino)ethyl)-2-(4-trifluoromethylphenyl)pyrid-5-ylmethylamine
<b>A32</b>	Int. A50		N-(2-(diethylamino)ethyl)-2-(4-chlorophenyl)pyrimid-5-ylmethylamine
<b>A33</b>	Int. A51		N-(2-(diethylamino)ethyl)-2-(4-trifluoromethylphenyl)pyrimid-5-ylmethylamine
<b>A34</b>	Int. A21		N-(2-(1-piperidino)ethyl)-4-(4-trifluoromethylphenyl)benzylamine
<b>A35</b>	Int. A22		(±)-N-(2-(diethylamino)ethyl)-1-(4-(4-chlorophenyl)phenyl)ethylamine
<b>A36</b>	Int. A54		N-(2-(diethylamino)ethyl)-3-(4-trifluoromethylphenoxy)benzylamine
<b>A37</b>	Int. A11		N-(2-(diethylamino)ethyl)-4-(4-trifluoromethylphenoxy)benzylamine
<b>A38</b>	Int. A14 Int. A21		tert-Butyl {2-[4-(4-trifluoromethylphenyl)-benzylamino]ethyl} ethylcarbamate
<b>A39</b>	Int. A16		N-(2-(diethylamino)ethyl)-3-(4-trifluoromethylbiphenyl-4-yl)propylamine
<b>A140</b>	Int. A55		N-(2-(diethylamino)ethyl)-2-(4-trifluoromethylbiphenyl-4-yloxy)ethylamine

A141	Int. A21 Int. A122		N-[(2-(diethylamino)-2-ethyl)propyl]-4-(4-trifluoromethylphenyl)benzylamine
A142	Int. A21 Int. A124		N-tert-Butylaminoethyl-4-(4-trifluoromethylphenyl)benzylamine

The following intermediates were made by the method of Intermediate A4:

No.	Precursor	Name
A40	Int. A8	5-Hydroxymethyl-2-(4-chlorophenyl)pyrimidine
A41	Int. A53	4-chloro-5-hydroxymethyl-2-(4-trifluoromethylphenyl)pyrimidine

The following intermediates were made by the method of Intermediate A5:

No.	Precursor	Name
A50	Int. A40	5-Formyl-2-(4-chlorophenyl)pyrimidine
A51	Int. A9	5-Formyl-2-(4-trifluoromethylphenyl)pyrimidine
A54	Int. A10	3-(4-trifluoromethylphenoxy)benzaldehyde

The following intermediate was made by the method of Intermediate A6:

No.	Precursors	Name
A52	diethyl ethoxymalonate, 4-trifluoromethylbenzamide.HCl	Ethyl 2-(4-trifluoromethylphenyl)-4-oxopyrimidine-5-carboxylate

The following intermediate was made by the method of Intermediate A7:

No.	Precursor	Name
A53	Int. A52	Ethyl 2-(4-trifluoromethylphenyl)-4-chloropyrimidine-5-carboxylate

5

The following intermediate was made by the method of Intermediate A16:

No.	Precursor	Name
A55	Int. A24	(4-trifluoromethylbiphenyl-4-yloxy)acetaldehyde

The following intermediates were made by the method of Intermediate A18, using Intermediate A17 and the appropriately substituted 1-alkyl-4-piperidone:

No.	Name
A60	N-(1-methylpiperidin-4-yl)-(4'-trifluoromethylphenyl)benzylamine
A61	N-(1-isopropylpiperidin-4-yl)-(4'-trifluoromethylphenyl)benzylamine
A62	N-(1-(2-methoxyethyl)piperidin-4-yl)-(4'-trifluoromethylphenyl)benzylamine

10

The following compounds are commercially available:

Intermediate B1, 2-thiouracil; Intermediate B2, 5-methyl-2-thiouracil ; Intermediate B3, 5-ethyl-2-thiouracil; Intermediate B4, 5-propyl-2-thiouracil; Intermediate B5, 5,6-dimethyl-2-thiouracil;

The following compounds are available by literature methods:

Intermediate B6, 5-carbethoxy-2-thiouracil (J. Amer. Chem. Soc. 794, 64 (1942));

5 Intermediate B7, 5,6-trimethylene-2-thiouracil (J. Amer. Chem. Soc. 3108, 81 (1959));


Intermediate B8, 5,6-tetramethylene-2-thiouracil (J. Org. Chem. 133, 18 (1953));

Intermediate B9, 5-methoxy-2-thiouracil (J. Chem. Soc. 4590 (1960)).

#### Intermediate B10 — 5-(2-hydroxyethyl)-2-thiouracil


10 A solution of ethyl formate (33.1 ml, 2.1 equiv) and  $\gamma$ -butyrolactone (15 ml, 1 equiv) in ether (400 ml) was added dropwise with stirring to a solution of potassium t-butoxide (52.5 g, 2.4 equiv) in tetrahydrofuran (400 ml). The mixture was allowed to warm to room temperature, and stirred overnight. The solvent was removed in vacuo, 2-propanol (600 ml) and thiourea (29.7 g, 2 equiv) were added, and the mixture was heated to reflux for 5h. After cooling to room temperature, the precipitate was filtered off, dissolved in water (500 ml), and washed twice with ether. The aqueous solution was acidified to  
15 pH5.5 with acetic acid, and the resulting precipitate was filtered off, washed thoroughly with water, and dried in vacuo; yield 23.85 g.  $^1\text{H-NMR}$  ( $d_6$ -DMSO)  $\delta$  2.36 (2H,t), 3.47 (2H,m), 4.57 (1H,m), 7.24 (1H,s), 12.2 & 12.4 (each 1H, br s); MS (APCI-) found (M-H) = 171;  $\text{C}_6\text{H}_8\text{N}_2\text{O}_2\text{S}$  requires 172.

#### Intermediate B111 — Ethyl (2,4-dioxo-4 *H*-benzo[d][1,3]oxazin-1-yl)acetate




20 Isatoic anhydride (10g, 1 equiv) in dimethylformamide (30ml) was added dropwise to a suspension of sodium hydride (2.45g, 60% in mineral oil, 1 equiv) in dimethylformamide (70ml) at room temperature. The reaction was stirred for 1 hour prior to the addition of ethyl bromoacetate (6.8ml, 1 equiv) and the resulting mixture stirred for 16 hours. Solvent evaporated, the residue suspended in water and the solid collected. The title compound was obtained by crystallisation from ethyl acetate (10.5g).  $^1\text{H-NMR}$   
25 ( $\text{CDCl}_3$ )  $\delta$  1.29 (3H,t), 4.27 (2H,q), 4.82 (2H,s), 6.96 (1H,d), 7.33 (1H,t), 7.74 (1H, dt), 8.19 (1H,dd).


#### Intermediate B112 — Ethyl (4-oxo-2-thioxo-3,4-dihydro-2 *H*-quinazolin-1-yl)acetate



30 Intermediate B111 (2.64g, 1equiv) and thiourea (2.42g, 4 equiv) in 1-methyl-2-pyrrolidinone (40ml) was heated to 180°C for 2 hours. After cooling the mixture was treated with water and the resultant solid collected by filtration. This solid was applied to a column (silica, 2% methanol/dichloromethane) to afford the title compound as a colourless solid (0.169g).  $^1\text{H-NMR}$  ( $\text{CDCl}_3$ )  $\delta$  1.22 (3H,t), 4.21 (2H,q), 5.53 (2H,br s), 7.46 (1H,t), 7.53 (1H,d), 7.81 (1H,dt), 8.07 (1H,dd).

**Intermediate B113 — Methyl 3-[3-(1-phenylmethanoyl)thioureido]thiophene-2-carboxylate**

Methyl-3-amino-2-thiophene carboxylate (30g, 1 equiv) and benzoyl isothiocyanate (46ml, 1.8 equiv) in acetone (250ml) were heated to 65°C for 30 minutes. After cooling the solution was concentrated and the resulting solid filtered off and dried (40.54g). <sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ 3.98 (3H,s), 7.54 (4H,m), 7.94 (2H,m), 8.81 (1H,d), 9.15 (1H,br s); MS (APCI+) found (M+1) = 321; C<sub>14</sub>H<sub>12</sub>N<sub>2</sub>O<sub>3</sub>S<sub>2</sub> requires 320.

**Intermediate B114 — 2-Thioxo-2,3-dihydro-1 H-thieno[3,2-d]pyrimidin-4-one**

Potassium hydroxide (13.83g, 2 equiv) was dissolved in ethanol (1000ml) and then poured onto intermediate B113 (40.54g, 1 equiv) with stirring. The mixture was heated to reflux for 1 hour and after cooling the title compound was obtained by filtration (17.32g). <sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ 6.87 (1H,d), 7.77 (1H,d), 10.46 (2H,br s); MS (APCI-) found (M-1) = 183; C<sub>6</sub>H<sub>4</sub>N<sub>2</sub>OS<sub>2</sub> requires 184.

The following intermediates were prepared by the method of Intermediate B10

No.	Precursor	Name
B11	monoethyl succinate	5-carboxymethyl-2-thiouracil
B12	ethyl ethoxyacetate	5-ethoxy-2-thiouracil
B13	ethyl (methylthio)acetate	5-methylthio-2-thiouracil

**Intermediate B20 — 2-(4-fluorobenzylthio)-5-methylpyrimidin-4-one**

A mixture of Intermediate B2 (9.45 g, 1 equiv), 4-fluorobenzyl chloride (7.96 ml, 1 equiv), potassium carbonate (18.4 g, 2 equiv) and dimethyl formamide (100 ml) was stirred at 90°C under argon for 16h. The DMF was removed in vacuo, water was added, and the product was extracted into ethyl acetate. The organic layer was dried and evaporated, and the residue was triturated with petroleum ether to obtain the title compound as a white solid (8.76 g). <sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ 2.02 (3H,s), 4.38 (2H,s), 6.97 (2H,m), 7.35 (2H,m), 7.74 (1H,s); MS (APCI+) found (M+1) = 251; C<sub>12</sub>H<sub>11</sub>FN<sub>2</sub>OS requires 250.

The following intermediates were prepared by the method of Intermediate B20:

No.	Precursor	Name
B21	Int. B1	2-(4-fluorobenzylthio)pyrimidin-4-one
B22	Int. B3	2-(4-fluorobenzylthio)-5-ethylpyrimidin-4-one
B23	Int. B4	2-(4-fluorobenzylthio)-5-propylpyrimidin-4-one
B24	Int. B6	2-(4-fluorobenzylthio)-5-ethoxycarbonylpyrimidin-4-one
B25	Int. B10	2-(4-fluorobenzylthio)-5-(2-hydroxyethyl)pyrimidin-4-one
B26	Int. B5	2-(4-fluorobenzylthio)-5,6-dimethylpyrimidin-4-one



B27	Int. B7	2-(4-fluorobenzylthio)-5,6-trimethylenepyrimidin-4-one
B28	Int. B8	2-(4-fluorobenzylthio)-5,6-tetramethylenepyrimidin-4-one
B29	Int. B9	2-(4-fluorobenzylthio)-5-methoxypyrimidin-4-one
B30	Int. B12	2-(4-fluorobenzylthio)-5-ethoxypyrimidin-4-one
B31	Int. B13	2-(4-fluorobenzylthio)-5-methylthiopyrimidin-4-one
B132	Int. B114	2-(4-fluorobenzylthio)-1 <i>H</i> -thieno[3,2- <i>d</i> ]pyrimidin-4-one

The following intermediates were prepared by method of Intermediate B20 and the appropriate benzyl chloride.

No.	Precursor	Name
B133	Int. B7 2,3-difluorobenzyl chloride	2-(2,3-difluorobenzylthio)-5,6-trimethylenepyrimidin-4-one
B134	Int. B7 3,4-difluorobenzyl chloride	2-(3,4-difluorobenzylthio)-5,6-trimethylenepyrimidin-4-one
B135	Int. B7 2,3,4-trifluorobenzyl chloride	2-(2,3,4-trifluorobenzylthio)-5,6-trimethylenepyrimidin-4-one
B136	Int. B7 2-fluorobenzyl chloride	2-(2-fluorobenzylthio)-5,6-trimethylenepyrimidin-4-one

**Intermediate B37 — 2-(4-fluorobenzylthio)-5-hydroxymethylpyrimidin-4-one**

- 5 Borane-tetrahydrofuran complex (143 ml, 2.2 equiv, 1.0M in THF) was added dropwise to an ice-cooled solution of Intermediate B24 (20 g, 1 equiv) in dry THF (700 ml) under argon with stirring. After a further 30 min at 0°C, the mixture was allowed to warm to room temperature and stirring continued overnight. The solvent was evaporated, 50% aqueous acetic acid (500 ml) was added with stirring, and the mixture was evaporated to dryness. The residue was digested with hot water (500 ml) for 5 min; then
- 10 the solid was filtered off. Both this solid and the filtrate were extracted with dichloromethane, and the organic extracts were combined and purified by chromatography (silica, 2-8% methanol in dichloromethane). Product fractions were evaporated to a white solid (6.14 g). <sup>1</sup>H-NMR (d<sub>6</sub>-DMSO) δ 4.25 (2H,s), 4.39 (2H,s), 7.14 (2H,t), 7.45 (2H,m), 7.82 (1H, br s); MS (APCI+) found (M+1) = 267; C<sub>12</sub>H<sub>11</sub>FN<sub>2</sub>O<sub>2</sub>S requires 266.

15 **Intermediate B38 — 2-(4-fluorobenzylthio)-5-isopropoxycarbonylmethylpyrimidin-4-one**

- A mixture of Intermediate B11 (2.60 g, 1 equiv), 4-fluorobenzyl bromide (1.74 ml, 1 equiv) and 2-propanol (50 ml) was stirred at reflux for 3h, then concentrated to a slurry in vacuo and diluted with ether. The solid was filtered off, washed with ether and dried; yield 2.87 g. <sup>1</sup>H-NMR (d<sub>6</sub>-DMSO) δ 1.17 (6H,d), 3.31 (2H,s), 4.40 (2H,s), 4.89 (1H,m), 7.14 (2H,t), 7.45 (2H,m), 7.84 (1H,s); MS (APCI+) found (M+1) = 325; C<sub>15</sub>H<sub>17</sub>FN<sub>2</sub>O<sub>3</sub>S requires 324.
- 20

**Intermediate B40 — 1-(tert-Butoxycarbonylmethyl)-2-(4-fluorobenzylthio)-5-methylpyrimidin-4-one**

A mixture of Intermediate B20 (6.30 g, 1 equiv), t-butyl iodoacetate (6.1 g, 1 equiv), diisopropylethylamine (5.27 ml, 1.2 equiv) and dichloromethane (100 ml) was stirred at ambient temperature under

argon for 16h, then the solution was washed with aq. ammonium chloride and aq. sodium bicarbonate, dried and evaporated. Chromatography (silica, ethyl acetate + 0.5% v/v aq. ammonia) followed by crystallisation from ethyl acetate gave the title compound as a white solid (3.36 g). <sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ 1.44 (9H,s), 2.01 (3H,d), 4.36 (2H,s), 4.51 (2H,s), 6.98 (3H,m), 7.36 (2H,m); MS (APCI+) found (M+1) = 365; C<sub>18</sub>H<sub>21</sub>FN<sub>2</sub>O<sub>3</sub>S requires 364.

The following intermediates were prepared by the method of Intermediate B40:

No.	Precursor	Name
B41	Int. B21	1-(tert-Butoxycarbonylmethyl)-2-(4-fluorobenzylthio)pyrimidin-4-one
B42	Int. B22	1-(tert-Butoxycarbonylmethyl)-2-(4-fluorobenzylthio)-5-ethylpyrimidin-4-one
B43	Int. B23	1-(tert-Butoxycarbonylmethyl)-2-(4-fluorobenzylthio)-5-propylpyrimidin-4-one
B44	Int. B24	1-(tert-Butoxycarbonylmethyl)-2-(4-fluorobenzylthio)-5-ethoxycarbonylpyrimidin-4-one
B45	Int. B38	1-(tert-Butoxycarbonylmethyl)-2-(4-fluorobenzylthio)-5-isopropoxycarbonylmethylpyrimidin-4-one
B46	Int. B37	1-(tert-Butoxycarbonylmethyl)-2-(4-fluorobenzylthio)-5-hydroxymethylpyrimidin-4-one
B47	Int. B25	1-(tert-Butoxycarbonylmethyl)-2-(4-fluorobenzylthio)-5-(2-hydroxyethyl)pyrimidin-4-one
B48	Int. B26	1-(tert-Butoxycarbonylmethyl)-2-(4-fluorobenzylthio)-5,6-dimethylpyrimidin-4-one
B49	Int. B27	1-(tert-Butoxycarbonylmethyl)-2-(4-fluorobenzylthio)-5,6-trimethylene-pyrimidin-4-one
B50	Int. B28	1-(tert-Butoxycarbonylmethyl)-2-(4-fluorobenzylthio)-5,6-tetramethylene-pyrimidin-4-one
B51	Int. B29	1-(tert-Butoxycarbonylmethyl)-2-(4-fluorobenzylthio)-5-methoxy-pyrimidin-4-one
B52	Int. B30	1-(tert-Butoxycarbonylmethyl)-2-(4-fluorobenzylthio)-5-ethoxypyrimidin-4-one
B53	Int. B31	1-(tert-Butoxycarbonylmethyl)-2-(4-fluorobenzylthio)-5-methylthio-pyrimidin-4-one
B154	Int. B133	1-(tert-Butoxycarbonylmethyl)-2-(2,3-difluorobenzylthio)-5,6-tetramethylenepyrimidin-4-one
B155	Int. B134	1-(tert-Butoxycarbonylmethyl)-2-(3,4-difluorobenzylthio)-5,6-tetramethylenepyrimidin-4-one
B156	Int. B135	1-(tert-Butoxycarbonylmethyl)-2-(2,3,4-trifluorobenzylthio)-5,6-tetramethylenepyrimidin-4-one
B157	Int. B136	1-(tert-Butoxycarbonylmethyl)-2-(2-fluorobenzylthio)-5,6-tetramethylenepyrimidin-4-one
B158	Int. B132	1-(tert-Butoxycarbonylmethyl)-2-(4-fluorobenzylthio)-4-oxo-4 H-thieno[3,2-d]pyrimidin-1-one


The following intermediate was prepared by the method of Intermediate B20:

No.	Precursor	Name
B159	B112	Ethyl [2-(4-fluorobenzylthio)-4-oxo-4 <i>H</i> -quinazolin-1-yl]acetate

**Intermediate B56 — 1-(tert-Butoxycarbonylmethyl)-2-(4-fluorobenzylthio)-5-chloropyrimidin-4-one**













- 5 A mixture of Intermediate B41 (7.45 g, 1 equiv), N-chlorosuccinimide (2.84 g, 1 equiv) and carbon tetrachloride (150 ml) was stirred at reflux under argon for 2h, then the solution was evaporated. Chromatography (silica, ethyl acetate) followed by trituration with ether gave the title compound as a white solid (4.45 g). <sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ 1.45 (9H,s), 4.40 (2H,s), 4.50 (2H,s), 6.99 (2H,m), 7.35 (2H,m), 7.40 (1H,s); MS (APCI+) found (M+1) = 385/387; C<sub>17</sub>H<sub>18</sub>ClFN<sub>2</sub>O<sub>3</sub>S requires 384/386.
- 10 **Intermediate B57 — 1-(tert-Butoxycarbonylmethyl)-2-(4-fluorobenzylthio)-5-bromopyrimidin-4-one**
- Prepared as Intermediate B56, but using N-bromosuccinimide in place of N-chlorosuccinimide. <sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ 1.45 (9H,s), 4.40 (2H,s), 4.49 (2H,s), 6.99 (2H,m), 7.35 (2H,m), 7.53 (1H,s); MS (APCI+) found (M+1) = 429/431; C<sub>17</sub>H<sub>18</sub>BrFN<sub>2</sub>O<sub>3</sub>S requires 428/430.
- 15 **Intermediate B58 — 1-(tert-Butoxycarbonylmethyl)-2-(4-fluorobenzylthio)-5-methylsulfinylpyrimidin-4-one**
- m*-Chloroperbenzoic acid (0.93 g) was added to an ice-cooled slurry of Intermediate B53 (1.50 g) in dichloromethane (20 ml). The resulting solution was allowed to warm to room temperature and stirred for 30 min, then washed with aq. sodium bicarbonate. Chromatography (silica, 3-8% methanol in ethyl acetate) gave the title compound as a white solid (1.15 g). <sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ 1.46 (9H,s), 2.94 (3H,s), 4.51 (4H,m), 7.01 (2H,m), 7.37 (2H,m), 7.60 (1H,s); MS (APCI+) found (M+1) = 413; C<sub>18</sub>H<sub>21</sub>FN<sub>2</sub>O<sub>4</sub>S<sub>2</sub> requires 412.
- 20











**Intermediate B60 — 1-(Carboxymethyl)-2-(4-fluorobenzylthio)-5-methylpyrimidin-4-one**




- 25 Intermediate B40 (3.88 g) was added to solution of trifluoroacetic acid (10 ml) in dichloromethane (20 ml) under argon, and stirred overnight at room temperature. Evaporation of the solvent and trituration with ether gave the title compound as a white solid (3.04 g). <sup>1</sup>H-NMR (d<sub>6</sub>-DMSO) δ 1.81 (3H,d), 4.42 (2H,s), 4.66 (2H,s), 7.14 (2H,m), 7.47 (2H,m), 7.63 (1H,m); MS (APCI+) found (M+1) = 309; C<sub>14</sub>H<sub>13</sub>FN<sub>2</sub>O<sub>3</sub>S requires 308.
- 30 The following intermediates were prepared by the method of Intermediate B60:

No.	Precursor	Structure	Name
-----	-----------	-----------	------

B61	Int. B41		1-(Carboxymethyl)-2-(4-fluorobenzylthio)-pyrimidin-4-one
B62	Int. B42		1-(Carboxymethyl)-2-(4-fluorobenzylthio)-5-ethylpyrimidin-4-one
B63	Int. B43		1-(Carboxymethyl)-2-(4-fluorobenzylthio)-5-propylpyrimidin-4-one
B64	Int. B44		1-(Carboxymethyl)-2-(4-fluorobenzylthio)-5-ethoxycarbonylpyrimidin-4-one
B65	Int. B45		1-(Carboxymethyl)-2-(4-fluorobenzylthio)-5-isopropoxycarbonylmethylpyrimidin-4-one
B66	Int. B46		1-(Carboxymethyl)-2-(4-fluorobenzylthio)-5-hydroxymethylpyrimidin-4-one
B67	Int. B47		1-(Carboxymethyl)-2-(4-fluorobenzylthio)-5-(2-hydroxyethyl)pyrimidin-4-one
B68	Int. B48		1-(Carboxymethyl)-2-(4-fluorobenzylthio)-5,6-dimethylpyrimidin-4-one
B69	Int. B49		1-(Carboxymethyl)-2-(4-fluorobenzylthio)-5,6-trimethylenepyrimidin-4-one
B70	Int. B50		1-(Carboxymethyl)-2-(4-fluorobenzylthio)-5,6-tetramethylenepyrimidin-4-one
B71	Int. B56		1-(Carboxymethyl)-2-(4-fluorobenzylthio)-5-chloropyrimidin-4-one
B72	Int. B57		1-(Carboxymethyl)-2-(4-fluorobenzylthio)-5-bromopyrimidin-4-one

<b>B73</b>	<b>Int. B51</b>		1-(Carboxymethyl)-2-(4-fluorobenzylthio)-5-methoxypyrimidin-4-one
<b>B74</b>	<b>Int. B52</b>		1-(Carboxymethyl)-2-(4-fluorobenzylthio)-5-ethoxypyrimidin-4-one
<b>B75</b>	<b>Int. B53</b>		1-(Carboxymethyl)-2-(4-fluorobenzylthio)-5-methylthiopyrimidin-4-one
<b>B76</b>	<b>Int. B58</b>		1-(Carboxymethyl)-2-(4-fluorobenzylthio)-5-methylsulfinylpyrimidin-4-one
<b>B177</b>	<b>Int. B154</b>		1-(Carboxymethyl)-2-(2,3-difluorobenzylthio)-5,6-trimethylenepyrimidin-4-one
<b>B178</b>	<b>Int. B155</b>		1-(Carboxymethyl)-2-(3,4-difluorobenzylthio)-5,6-trimethylenepyrimidin-4-one
<b>B179</b>	<b>Int. B156</b>		1-(Carboxymethyl)-2-(2,3,4-trifluorobenzylthio)-5,6-trimethylenepyrimidin-4-one
<b>B180</b>	<b>Int. B157</b>		1-(Carboxymethyl)-2-(2-fluorobenzylthio)-5,6-trimethylenepyrimidin-4-one
<b>B181</b>	<b>Int. B158</b>		[2-(4-Fluorobenzylthio)-4-oxo-4 H-thieno[3,2-d]pyrimidin-1-yl]acetic acid
<b>B182</b>	<b>Int. B159</b>		[2-(4-Fluorobenzylthio)-4-oxo-4 H-quinazolin-1-yl]acetic acid

**Intermediate B80 — 1-(N-Methyl-N-(4-(4-chlorophenyl)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzylthio)-5-(2-azidoethyl)pyrimidin-4-one**




A mixture of Example 39 (1.88 g, 1 equiv), methanesulfonic anhydride (0.713 g, 1.2 equiv), triethylamine (0.665 ml) and dichloromethane (20 ml) was stirred at 0°C for 4h. The solution was washed with water, dried and evaporated to a pale foam (2.4 g). This was dissolved in dimethylformamide (20 ml), sodium azide (0.266 g, 1.2 equiv) was added, and the mixture was stirred under argon at room temperature overnight. The solvent was evaporated, the residue was partitioned between water and dichloromethane, and the organic layer was dried and evaporated. Chromatography (silica, ethyl acetate) gave the title compound as a white solid. <sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ 2.66 (2H,m), 2.88 (3H, s), 3.60(2H, m), 4.46-4.64 (6H, m), 6.84-7.50 (12H, m), 8.02 (1H,s); MS (APCI+) found (M+1) = 577/579; C<sub>29</sub>H<sub>26</sub>ClFN<sub>6</sub>O<sub>2</sub>S requires 576/578.

The following compound was prepared by the method of Intermediate B80


No.	Precursor	Name
B81	Example 42	1-(N-(2-(Diethylamino)ethyl)-N-(4-(4-trifluoromethylphenyl)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(2-azidoethyl)pyrimidin-4-one

**Example 1 — 1-(N-(2-(Diethylamino)ethyl)-N-(4-(4-trifluoromethylphenyl)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-ethylpyrimidin-4-one bitartrate**



A mixture of Intermediate A30 (0.403 g, 1 equiv), Intermediate B62 (0.371 g, 1 equiv), HATU (0.426 g, 1.2 equiv), di-isopropylethylamine (0.482 ml, 2.4 equiv) and dichloromethane (15 ml) was stirred at room temperature overnight, then washed with aqueous ammonium chloride and aqueous sodium bicarbonate. The organic layer was dried and evaporated, and the product purified by chromatography (silica, 5% methanol in dichloromethane). Product fractions were evaporated to a white foam (0.627 g). This free base (0.612 g) was dissolved in methanol (10 ml), tartaric acid (0.14 g) was added, the mixture was stirred for 5 mins then evaporated. Trituration with ether gave the bitartrate salt as a white solid (0.622 g). <sup>1</sup>H-NMR (d<sub>6</sub>-DMSO, ca 1:1 rotamer mixture) δ 0.96 (3H,m), 1.07 (6H,m), 2.27 (2H,m), 2.59 (2H,m), 2.84 (2H,m), 3.37/3.50 (4H,m), 4.26 (2H,s), 4.39/4.43 (2H,2x s), 4.64/4.72 (2H,2x s), 4.94/5.09 (2H,2x s), 7.11/7.14 (2H,2x m), 7.36-7.49 (5H, m), 7.63/7.72 (2H,2x d), 7.84 (4H,m); MS (APCI+) found (M+1) = 655; C<sub>35</sub>H<sub>38</sub>F<sub>4</sub>N<sub>4</sub>O<sub>2</sub>S requires 654.


**Example 2 — 1-(N-(2-(Diethylamino)ethyl)-N-(2-(4-trifluoromethylphenyl)pyrid-5-ylmethyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-ethylpyrimidin-4-one bitartrate**



Prepared from intermediates A31 and B62 by the method of Example 1. <sup>1</sup>H-NMR (d<sub>6</sub>-DMSO, ca 2:1 rotamer mixture) δ 0.93 (6H,m), 1.08 (3H,m), 2.27 (2H,m), 2.66 (4H,m), 3.39/3.45 (4H,m), 4.21 (2H,s), 4.39/4.42 (2H,2x s), 4.66/4.77 (2H,2x s), 4.97/5.10 (2H,2x s), 7.09/7.12 (2H,2x t), 7.42/7.49 (2H,2x t),

7.79/7.86 (1H, 2x dd), 7.87 (2H, d), 7.97/8.06 (1H, 2x dd), 8.28 (2H, d), 8.62/8.71 (2H, 2x s); MS (APCI+) found (M+1) = 656; C<sub>34</sub>H<sub>37</sub>F<sub>4</sub>N<sub>5</sub>O<sub>2</sub>S requires 655.

**Example 3(a) — 1-(N-(2-(Diethylamino)ethyl)-N-(4-(4-trifluoromethylphenyl)benzyl)-aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5,6-trimethylenepyrimidin-4-one**



Intermediate B69 (87.1g, 0.26 mol.) was suspended in dichloromethane (2.9 litre). 1-Hydroxybenzotriazole hydrate (35.2g, 0.26 mol.) and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (99.7g, 0.52 mol.) were added and the suspension stirred for 45 minutes by which time complete solution had been obtained. Intermediate A30 (91.2g, 0.26 mol.) was added as a solution in dichloromethane (100ml) over 5 minutes and the solution stirred for 4 hours. Saturated ammonium chloride solution:water mixture (1:1, 1 litre) was added and the solution stirred for 10 minutes. The organic phase was separated and extracted with saturated ammonium chloride:water mixture (1:1, 1 litre), extracts were pH 6. The organic phase was separated and extracted with water (1 litre) containing acetic acid (10ml), extract pH 5. The dichloromethane layer was separated and extracted with saturated sodium carbonate solution:water:saturated brine mixture (1:3:0.2, 1 litre), pH 10.5, then with saturated brine:water mixture (1:1, 1 litre). The brown solution was dried over anhydrous sodium sulfate in the presence of decolourising charcoal (35g), filtered and the solvent removed *in vacuo* to give a dark brown foam. The foam was dissolved in *iso*-propyl acetate (100ml) and the solvent removed *in vacuo*. The dark brown gummy residue was dissolved in boiling *iso*-propyl acetate (500ml), cooled to room temperature, seeded and stirred overnight. The pale cream solid produced was filtered off and washed with *iso*-propyl acetate (100ml). The solid was sucked dry in the sinter for 1 hour then recrystallized from *iso*-propyl acetate (400ml). After stirring overnight the solid formed was filtered off, washed with *iso*-propyl acetate (80ml) and dried *in vacuo* to give the title compound, 110g, 63.5% yield. <sup>1</sup>H NMR (CDCl<sub>3</sub>, ca 1.9:1 rotamer mixture) δ 0.99 (6H, t), 2.10 (2H, m), 2.50 (4H, q), 2.58/2.62 (2H, 2 x t), 2.70/2.82 (2H, 2 x t), 2.86 (2H, t), 3.28/3.58 (2H, 2 x t), 4.45/4.52 (2H, 2 x s), 4.68/4.70 (2H, 2 x s), 4.93 (2H, s), 6.95 (2H, m), 7.31 (2H, d), 7.31/7.37 (2H, 2 x m), 7.48/7.52 (2H, d), 7.65 (2H, m), 7.72 (2H, m); MS (APCI) (M+H)<sup>+</sup> 667; mp 125°C (by DSC - assymetric endotherm).


**Example 3(b) — 1-(N-(2-(Diethylamino)ethyl)-N-(4-(4-trifluoromethylphenyl)benzyl)-aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5,6-trimethylenepyrimidin-4-one bitartrate**

Prepared from intermediates A30 and B69 by the method of Example 1. <sup>1</sup>H-NMR (d<sub>6</sub>-DMSO, ca 1:1 rotamer mixture) δ 0.92/0.99 (6H, 2x t), 1.99 (2H, m), 2.54 (6H, m), 2.68/2.74 (4H, m), 3.36 (2H, m), 4.21 (2H, s), 4.37/4.44 (2H, 2x s), 4.63/4.74 (2H, 2x s), 4.89/5.13 (2H, 2x s), 7.08/7.14 (2H, 2x m), 7.36-7.50 (4H, m), 7.64/7.70 (2H, 2x d), 7.83 (4H, m); MS (APCI+) found (M+1) = 667; C<sub>36</sub>H<sub>38</sub>F<sub>4</sub>N<sub>4</sub>O<sub>2</sub>S requires 666.

**Example 3(c) — 1-(N-(2-(Diethylamino)ethyl)-N-(4-(4-trifluoromethylphenyl)benzyl)-aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5,6-trimethylenepyrimidin-4-one hydrochloride**


The free base from Example 3(a) (3.00g, 0.0045 mol) was suspended with stirring in isopropanol (30 ml) and warmed to 45°C to give a clear solution. The solution was then cooled to ambient temperature and conc. hydrochloric acid (0.40 ml, 0.045 mol) was added. The resultant slurry was then stirred at ambient temperature for 35 minutes, before being cooled to 0°C for 35 minutes. The slurry was then filtered and washed with isopropanol (10 ml), followed by heptane (30 ml), before being dried under vacuum to give the title compound as a white solid (3.00 g, 95%). <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 1.38 (6H, t), 2.08 (2H, m), 2.82 (2H, t), 2.99 (2H, t), 3.19 (4H, m), 3.35 (2H, m), 3.97 (2H, s), 4.42 (2H, s), 4.81 (2H, s), 4.99 (2H, s), 6.87 (2H, t), 7.26 (2H, t), 7.33 (2H, d), 7.41 (2H, d), 7.53 (2H, d), 7.71 (2H, d), 11.91 (1H, s).

**Example 4 — 1-(N-(2-(Diethylamino)ethyl)-N-(2-(4-trifluoromethylphenyl)pyrid-5-ylmethyl)-aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5,6-trimethylenepyrimidin-4-one bitartrate**




Prepared from intermediates A31 and B69 by the method of Example 1. <sup>1</sup>H-NMR (d<sub>6</sub>-DMSO, ca 3:1 rotamer mixture) δ 0.92/0.98 (6H,t), 1.99 (2H,m), 2.53 (6H,m), 2.68/2.75 (4H,m), 3.41 (2H,m), 4.22 (2H,s), 4.37/4.42 (2H,2x s), 4.66/4.79 (2H,2x s), 4.93/5.13 (2H,2x s), 7.07/7.12 (2H,2x t), 7.39/7.47 (2H,2x t), 7.77/7.86 (1H,2x dd), 7.87 (2H,d), 7.98/8.05 (1H,2x dd), 8.28 (2H,d), 8.61/8.69 (1H,2x s); MS (APCI+) found (M+1) = 668; C<sub>35</sub>H<sub>37</sub>F<sub>4</sub>N<sub>5</sub>O<sub>2</sub>S requires 667.

**Example 5 — 1-(N-(2-(Diethylamino)ethyl)-N-(2-(4-trifluoromethylphenyl)pyrimid-5-ylmethyl)-aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5,6-trimethylenepyrimidin-4-one bitartrate**



Prepared from intermediates A33 and B69 by the method of Example 1. <sup>1</sup>H-NMR (d<sub>6</sub>-DMSO, ca 3:1 rotamer mixture) δ 0.92/1.09 (6H,t), 1.96 (2H,m), 2.60 (6H,m), 2.75 (4H,m), 3.48 (2H,m), 4.23 (2H,s), 4.38/4.40 (2H,2x s), 4.65/4.81 (2H,2x s), 4.97/5.11 (2H,2x s), 7.07/7.10 (2H,2x t), 7.38/7.44 (2H,2x t), 7.91 (2H,d), 8.57 (2H,d), 8.84/8.93 (2H,2x s); MS (APCI+) found (M+1) = 669; C<sub>34</sub>H<sub>36</sub>F<sub>4</sub>N<sub>6</sub>O<sub>2</sub>S requires 668.

**Example 6 — 1-(N-Methyl-N-(2-(4-chlorophenyl)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(2-aminoethyl)pyrimidin-4-one hydrochloride**




A solution of Intermediate B80 (0.228 g) in ethanol (20 ml) was hydrogenated over 10% palladium on charcoal (0.09 g) at atmospheric pressure for 2 days. The catalyst was filtered off, the solvent was




removed in vacuo, and the resulting oil was purified by chromatography (silica, 10% methanolic ammonia in dichloromethane). The free base was dissolved in dichloromethane (5 ml), and an equimolar quantity of hydrogen chloride in ether added. The solvent was removed in vacuo, and the residue triturated with ether; yield 0.132 g). <sup>1</sup>H-NMR (d<sub>6</sub>-DMSO, ca 2:1 rotamer mixture) δ 2.58 (2H,m), 2.87/2.99 (3H,2x s), 2.99 (2H,m), 4.40/4.45 (2H,2x s), 4.57/4.66 (2H,2x s), 4.97/5.00 (2H,2x s), 7.16 (2H,m), 7.33/7.38 (2H,2x d), 7.4-7.7 (9H,m), 8.0 (2H,br m); MS (APCI+) found (M+1) = 551/553; C<sub>29</sub>H<sub>28</sub>ClFN<sub>4</sub>O<sub>2</sub>S requires 550/552.

**Example 7 — 1-(N-Methyl-N-(2-(4-chlorophenyl)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(2-acetamidoethyl)pyrimidin-4-one**



A solution of Example 6 (0.173 g, 1 equiv), acetic anhydride (0.033 ml, 1.1 equiv) and diisopropylamine (0.066 ml, 1.2 equiv) in dichloromethane (10 ml) was stirred at room temperature overnight. The solution was washed with aq. ammonium chloride and aq. sodium bicarbonate, then the organic layer was dried and evaporated. The residue was triturated with ether to obtain the title compound as a white solid (0.156 g). <sup>1</sup>H-NMR (CDCl<sub>3</sub>, ca 2:1 rotamer mixture) δ 1.96 (3H,s), 2.64 (2H,m), 2.96/3.10 (3H, 2x s), 3.49 (2H,m), 4.46-4.64 (6H,m), 6.77 (1H,br t), 6.97-7.16 (3H,m), 7.26-7.49 (10H,m); MS (APCI+) found (M+1) = 593/595; C<sub>31</sub>H<sub>30</sub>ClFN<sub>4</sub>O<sub>3</sub>S requires 592/594.

**Example 8 — 1-(N-(2-(Diethylamino)ethyl)-N-(2-(4-chlorophenyl)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(dimethylaminomethyl)pyrimidin-4-one**



Methanesulfonic anhydride (0.134 g, 1.2 equiv) was added to a solution of Example 37 (0.40 g, 1 equiv) and triethylamine (0.124 ml, 1.4 equiv) in dichloromethane (5 ml) at 0°C, then stirred at this temperature for 4 hours. The mixture was washed with water, dried and evaporated to yield the mesylate as a pale yellow solid. This was dissolved in a 2M solution of dimethylamine in THF (10 ml) and stirred at room temperature for 16 hours. The solvent and excess dimethylamine was removed in vacuo, and the product was purified by chromatography (silica, 5-20% methanol in ethyl acetate, then 1-10% methanolic ammonia in dichloromethane) to obtain the title compound. <sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ 0.98 (6H,t), 2.28/2.30 (each 3H,s), 2.46-2.65 (6H,m), 3.26/3.56 (2H,2x t), 3.33/3.36 (2H,2x s), 4.46/4.53/5.54/4.90 (4H,4x s), 4.67 (2H,s), 6.98 (2H,m), 7.21-7.50 (11H,m); MS (APCI+) found (M+1) = 650/652; C<sub>35</sub>H<sub>41</sub>ClFN<sub>5</sub>O<sub>2</sub>S requires 649/651.

The following Examples were made by the method of Example 1 except that in a few cases EDC (2 equiv) and 1-hydroxybenzotriazole (1 equiv) were used in place of HATU and di-isopropylamine, in an essentially similar procedure. Where indicated, the salts were subsequently prepared by the methods of Examples 1 or 6 as appropriate:

Ex. No.	Precursors	Structure	Name
20	Int. A3 Int. B60		1-(N-(2-(Diethylamino)ethyl)-N-(4-(4-chlorophenyl)benzyl)-aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-methylpyrimidin-4-one hydrochloride
21	Int. A26 Int. B60		1-(N-methyl-N-(2-(4-trifluoromethylphenyl)pyrid-5-ylmethyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-methylpyrimidin-4-one bitartrate
22	Int. A30 Int. B60		1-(N-(2-(Diethylamino)ethyl)-N-(4-(4-trifluoromethylphenyl)benzyl)-aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-methylpyrimidin-4-one bitartrate
23	Int. A31 Int. B60		1-(N-(2-(Diethylamino)ethyl)-N-(2-(4-trifluoromethylphenyl)pyrid-5-yl-methyl)-aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-methylpyrimidin-4-one bitartrate
24	Int. A32 Int. B60		1-(N-(2-(Diethylamino)ethyl)-N-(2-(4-chlorophenyl)pyrimid-5-yl-methyl)-aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-methylpyrimidin-4-one bitartrate
25	Int. A33 Int. B60		1-(N-(2-(Diethylamino)ethyl)-N-(2-(4-trifluoromethylphenyl)pyrimid-5-yl-methyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-methylpyrimidin-4-one bitartrate
26	Int. A35 Int. B60		(±)-1-(N-(2-(Diethylamino)ethyl)-N-(1-(4-(4-chlorophenyl)phenyl)ethyl)-aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-methylpyrimidin-4-one bitartrate
27	Int. A34 Int. B60		1-(N-(2-(1-piperidino)ethyl)-N-(4-(4-trifluoromethylphenyl)benzyl)-aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-methylpyrimidin-4-one bitartrate
28	Int. A25 Int. B62		1-(N-methyl-N-(4-(4-trifluoromethylphenyl)benzyl)-aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-ethylpyrimidin-4-one

29	Int. A3 Int. B62		1-(N-(2-(Diethylamino)ethyl)-N-(4-(4-chlorophenyl)benzyl)-aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-ethylpyrimidin-4-one bitartrate
30	Int. A26 Int. B62		1-(N-methyl-N-(2-(4-trifluoromethylphenyl)pyrid-5-yl-methyl)-aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-ethylpyrimidin-4-one
31	Int. A32 Int. B62		1-(N-(2-(Diethylamino)ethyl)-N-(2-(4-chlorophenyl)pyrimid-5-yl-methyl)-aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-ethylpyrimidin-4-one bitartrate
32	Int. A33 Int. B62		1-(N-(2-(Diethylamino)ethyl)-N-(2-(4-trifluoromethylphenyl)pyrimid-5-yl-methyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-ethylpyrimidin-4-one bitartrate
33	Int. A3 Int. B63		1-(N-(2-(Diethylamino)ethyl)-N-(4-(4-chlorophenyl)benzyl)-aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-propylpyrimidin-4-one bitartrate
34	Int. A30 Int. B63		1-(N-(2-(Diethylamino)ethyl)-N-(4-(4-trifluoromethylphenyl)benzyl)-aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-propylpyrimidin-4-one bitartrate
35	Int. A30 Int. B64		1-(N-(2-(Diethylamino)ethyl)-N-(4-(4-trifluoromethylphenyl)benzyl)-aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-ethoxycarbonylmethylpyrimidin-4-one bitartrate
36	Int. A30 Int. B65		1-(N-(2-(Diethylamino)ethyl)-N-(4-(4-trifluoromethylphenyl)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-isopropoxycarbonylmethylpyrimidin-4-one bitartrate
37	Int. A3 Int. B66		1-(N-(2-(Diethylamino)ethyl)-N-(4-(4-chlorophenyl)benzyl)-aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-hydroxymethylpyrimidin-4-one bitartrate

38	Int. A30 Int. B66		1-(N-(2-(Diethylamino)ethyl)-N-(4-(4-trifluoromethylphenyl)benzyl)-aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-hydroxymethylpyrimidin-4-one bitartrate
39	Int. A2 Int. B67		1-(N-methyl-N-(4-(4-chlorophenyl)benzyl)-aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(2-hydroxyethyl)pyrimidin-4-one bitartrate
40	Int. A3 Int. B67		1-(N-(2-(Diethylamino)ethyl)-N-(4-(4-chlorophenyl)benzyl)-aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(2-hydroxyethyl)pyrimidin-4-one bitartrate
41	Int. A31 Int. B67		1-(N-(2-(Diethylamino)ethyl)-N-(2-(4-trifluoromethylphenyl)pyridin-5-ylmethyl)-aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(2-hydroxyethyl)pyrimidin-4-one bitartrate
42	Int. A30 Int. B67		1-(N-(2-(Diethylamino)ethyl)-N-(4-(4-trifluoromethylphenyl)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(2-hydroxyethyl)pyrimidin-4-one bitartrate
43	Int. A30 Int. B68		1-(N-(2-(Diethylamino)ethyl)-N-(4-(4-trifluoromethylphenyl)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5,6-dimethylpyrimidin-4-one bitartrate
44	Int. A3 Int. B69		1-(N-(2-(Diethylamino)ethyl)-N-(4-(4-chlorophenyl)benzyl)-aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5,6-trimethylenepyrimidin-4-one bitartrate
45	Int. A3 Int. B70		1-(N-(2-(Diethylamino)ethyl)-N-(4-(4-chlorophenyl)benzyl)-aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5,6-tetramethylenepyrimidin-4-one bitartrate
46	Int. A30 Int. B70		1-(N-(2-(Diethylamino)ethyl)-N-(4-(4-trifluoromethylphenyl)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5,6-tetramethylenepyrimidin-4-one bitartrate

47	Int. A31 Int. B70		1-(N-(2-(Diethylamino)ethyl)-N-(2-(4-trifluoromethylphenyl)pyrid-5-yl-methyl)-aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5,6-tetramethylenepyrimidin-4-one bitartrate
49	Int. A30 Int. B71		1-(N-(2-(Diethylamino)ethyl)-N-(4-(4-trifluoromethylphenyl)benzyl)amino-carbonyl-methyl)-2-(4-fluorobenzyl)thio-5-chloropyrimidin-4-one bitartrate
50	Int. A3 Int. B71		1-(N-(2-(Diethylamino)ethyl)-N-(4-(4-chlorophenyl)benzyl)-aminocarbonyl-methyl)-2-(4-fluorobenzyl)thio-5-chloro-pyrimidin-4-one bitartrate
51	Int. A31 Int. B71		1-(N-(2-(Diethylamino)ethyl)-N-(2-(4-trifluoromethylphenyl)pyrid-5-yl-methyl)-aminocarbonyl-methyl)-2-(4-fluorobenzyl)thio-5-chloropyrimidin-4-one bitartrate
52	Int. A30 Int. B72		1-(N-(2-(Diethylamino)ethyl)-N-(4-(4-trifluoromethylphenyl)benzyl)amino-carbonyl-methyl)-2-(4-fluorobenzyl)thio-5-bromopyrimidin-4-one bitartrate
53	Int. A3 Int. B72		1-(N-(2-(Diethylamino)ethyl)-N-(4-(4-chlorophenyl)benzyl)-aminocarbonyl-methyl)-2-(4-fluorobenzyl)thio-5-bromo-pyrimidin-4-one bitartrate
54	Int. A30 Int. B73		1-(N-(2-(Diethylamino)ethyl)-N-(4-(4-trifluoromethylphenyl)benzyl)amino-carbonylmethyl)-2-(4-fluorobenzyl)thio-5-methoxypyrimidin-4-one bitartrate
55	Int. A31 Int. B73		1-(N-(2-(Diethylamino)ethyl)-N-(2-(4-trifluoromethylphenyl)pyrid-5-yl-methyl)-aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-methoxypyrimidin-4-one bitartrate
56	Int. A30 Int. B74		1-(N-(2-(Diethylamino)ethyl)-N-(4-(4-trifluoromethylphenyl)benzyl)amino-carbonylmethyl)-2-(4-fluorobenzyl)thio-5-ethoxypyrimidin-4-one bitartrate
57	Int. A31 Int. B74		1-(N-(2-(Diethylamino)ethyl)-N-(2-(4-trifluoromethylphenyl)pyrid-5-yl-methyl)-aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-ethoxypyrimidin-4-one bitartrate

58	Int. A31 Int. B75		1-(N-(2-(Diethylamino)ethyl)-N-(2-(4-trifluoromethylphenyl)pyrid-5-yl-methyl)-aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-methylthiopyrimidin-4-one bitartrate
59	Int. A30 Int. B75		1-(N-(2-(Diethylamino)ethyl)-N-(4-(4-trifluoromethylphenyl)benzyl)amino-carbonylmethyl)-2-(4-fluorobenzyl)thio-5-methylthiopyrimidin-4-one bitartrate
60	Int. A30 Int. B76		1-(N-(2-(Diethylamino)ethyl)-N-(4-(4-trifluoromethylphenyl)benzyl)amino-carbonylmethyl)-2-(4-fluorobenzyl)thio-5-methylsulfinylpyrimidin-4-one bitartrate
61	Int. A31 Int. B76		1-(N-(2-(Diethylamino)ethyl)-N-(2-(4-trifluoromethylphenyl)pyrid-5-yl-methyl)-aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-methylsulfinylpyrimidin-4-one bitartrate
62	Int. A30 Int. B177		1-(N-(2-(Diethylamino)ethyl)-N-(4-(4-trifluoromethylphenyl)benzyl)amino-carbonylmethyl)-2-(2,3-difluorobenzyl)-thio-5,6-trimethylenepyrimidin-4-one bitartrate
63	Int. A30 Int. B178		1-(N-(2-(Diethylamino)ethyl)-N-(4-(4-trifluoromethylphenyl)benzyl)amino-carbonylmethyl)-2-(3,4-difluorobenzyl)-thio-5,6-trimethylenepyrimidin-4-one bitartrate
64	Int. A30 Int. B179		1-(N-(2-(Diethylamino)ethyl)-N-(4-(4-trifluoromethylphenyl)benzyl)amino-carbonylmethyl)-2-(2,3,4-trifluorobenzyl)-thio-5,6-trimethylenepyrimidin-4-one bitartrate
65	Int. A30 Int. B180		1-(N-(2-(Diethylamino)ethyl)-N-(4-(4-trifluoromethylphenyl)benzyl)amino-carbonylmethyl)-2-(2-fluorobenzyl)thio-5,6-trimethylenepyrimidin-4-one bitartrate
66	Int. A25 Int. B69		1-(N-methyl-N-(4-(4-trifluoromethylphenyl)benzyl)-aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5,6-trimethylene-pyrimidin-4-one

67	Int. A34 Int. B69		1-(N-(2-(1-piperidino)ethyl)-N-(4-(4-trifluoromethylphenyl)benzyl)-aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5,6-trimethylenepyrimidin-4-one bitartrate
68	Int. A36 Int. B69		1-(N-(2-(Diethylamino)ethyl)-N-(3-(4-trifluoromethylphenoxy)benzyl)-aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5,6-trimethylenepyrimidin-4-one
69	Int. A37 Int. B69		1-(N-(2-(Diethylamino)ethyl)-N-(4-(4-trifluoromethylphenoxy)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5,6-trimethylenepyrimidin-4-one bitartrate
70	Int. A39 Int. B69		1-(N-(2-(Diethylamino)ethyl)-N-(4-(4-trifluoromethyl-biphenyl-4-yl)propyl)-aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5,6-trimethylenepyrimidin-4-one
71	Int. A39 Int. B62		1-(N-(2-(Diethylamino)ethyl)-N-(4-(4-trifluoromethyl-biphenyl-4-yl)propyl)-aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-ethylpyrimidin-4-one
72	Int. A140 Int. B62		1-(N-(2-(Diethylamino)ethyl)-N-(4-(4-trifluoromethyl-biphenyl-4-yloxy)ethyl)-aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-ethylpyrimidin-4-one
73	Int. A18 Int. B69		1-(N-(1-Ethyl-piperidin-4-yl)-N-(4-(4-trifluoromethylphenyl)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5,6-trimethylenepyrimidin-4-one bitartrate
74	Int. A141 Int. B69		1-(N-(2-Ethylamino-2-methylpropyl)-N-(4-(4-trifluoromethylphenyl)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5,6-trimethylenepyrimidin-4-one bitartrate
75	Int. A142 Int. B69		N-(2-tert-butylaminoethyl)-N-(4-(4-trifluoromethylphenyl)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5,6-trimethylenepyrimidin-4-one bitartrate

76	Int. A30 Int. B181		N-(2-Diethylaminoethyl)-2-[2-(4-fluorobenzylthio)-4-oxo-4 <i>H</i> -thieno[3,2- <i>d</i> ]pyrimidin-1-yl]-N-(4'-trifluoromethyl-biphenyl-4-ylmethyl)-acetamide bitartrate
77	Int. A30 Int. B182		N-(2-Diethylaminoethyl)-2-[2-(4-fluorobenzylthio)-4-oxo-4 <i>H</i> -quinazolin-1-yl]-N-(4'-trifluoromethyl-biphenyl-4-ylmethyl)-acetamide bitartrate
78	Int. A38 Int. B69		Ethyl-{2-[[2-(4-fluorobenzylthio)-4-oxo-4,5,6,7-tetrahydrocyclopentapyrimidin-1-yl]-ethanoyl]-{4'-trifluoromethylbiphenyl-4-ylmethyl}-amino}-ethyl} carbamic acid tert-butyl ester
79	Int. A60 Int. B69		1-(N-(1-Methylpiperidin-4-yl)-N-(4-(4-trifluoromethylphenyl)benzyl)amino-carbonylmethyl)-2-(4-fluorobenzyl)thio-5,6-trimethylenepyrimidin-4-one bitartrate
80	Int. A61 Int. B69		1-(N-(1-Isopropylpiperidin-4-yl)-N-(4-(4-trifluoromethylphenyl)benzyl)amino-carbonylmethyl)-2-(4-fluorobenzyl)thio-5,6-trimethylenepyrimidin-4-one bitartrate
81	Int. A62 Int. B69		1-(N-(1-(2-Methoxyethyl)piperidin-4-yl)-N-(4-(4-trifluoromethylphenyl)benzyl)-aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5,6-trimethylenepyrimidin-4-one bitartrate

The following compound was prepared by the method of Example 6:

No.	Precursor	Structure	Name
85	Int. B81		1-(N-(2-(Diethylamino)ethyl)-N-(4-(4-trifluoromethylphenyl)benzyl)amino-carbonylmethyl)-2-(4-fluorobenzyl)thio-5-(2-aminoethyl)pyrimidin-4-one bitartrate

The following compounds were prepared by the method of Example 7:

No.	Precursors	Structure	Name
-----	------------	-----------	------




90	Example 85, acetic anhydride		1-(N-(2-(Diethylamino)ethyl)-N-(4-(4-trifluoromethylphenyl)benzyl)amino-carbonylmethyl)-2-(4-fluorobenzyl)thio-5-(2-acetamidoethyl)pyrimidin-4-one bitartrate
91	Example 85, methane-sulfonic anhydride		1-(N-(2-(Diethylamino)ethyl)-N-(4-(4-trifluoromethylphenyl)benzyl)amino-carbonylmethyl)-2-(4-fluorobenzyl)thio-5-(2-methanesulfonamidoethyl)pyrimidin-4-one bitartrate
92	Example 85, methoxy-acetyl chloride		1-(N-(2-(Diethylamino)ethyl)-N-(4-(4-trifluoromethylphenyl)benzyl)amino-carbonylmethyl)-2-(4-fluorobenzyl)thio-5-(2-(methoxyacetamido)ethyl)pyrimidin-4-one bitartrate

The following example was prepared by the method of Intermediate B60. The salt was prepared by the method of example 1:

No.	Precursor	Structure	Name
93	Example 78		1-(N-(2-(Ethylamino)ethyl)-N-(4-(4-trifluoromethylphenyl)benzyl)-aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5,6-trimethylene-pyrimidin-4-one bitartrate

**Biological Data****1. Screen for Lp-PLA<sub>2</sub> inhibition.**

- 5 Enzyme activity was determined by measuring the rate of turnover of the artificial substrate (A) at 37 °C in 50mM HEPES (N-2-hydroxyethylpiperazine-N'-2-ethanesulphonic acid) buffer containing 150mM NaCl, pH 7.4.



(A)

Assays were performed in 96 well titre plates.

- 15 Recombinant Lp-PLA<sub>2</sub> was purified to homogeneity from baculovirus infected Sf9 cells, using a zinc chelating column, blue sepharose affinity chromatography and an anion exchange column. Following purification and ultrafiltration, the enzyme was stored at 6mg/ml at 4°C. Assay plates of compound or vehicle plus buffer were set up using automated robotics to a volume of 170µl. The reaction was initiated by the addition of 20µl of 10x substrate (A) to give a final substrate concentration of 20µM and 10 µl of diluted enzyme to a final 0.2nM Lp-PLA<sub>2</sub>.


- 20 The reaction was followed at 405 nm and 37 °C for 20 minutes using a plate reader with automatic mixing. The rate of reaction was measured as the rate of change of absorbance.

**Results**

- 25 The compounds described in the Examples were tested as described above and had IC<sub>50</sub> values in the range <0.1nM to 10 µM.

## Claims

1. A compound of formula (I):



(I)

in which:

$R^a$  is hydrogen, halogen,  $C_{(1-3)}$ alkyl,  $C_{(1-3)}$ alkoxy, hydroxy $C_{(1-3)}$ alkyl,  $C_{(1-3)}$ alkylthio,  $C_{(1-3)}$ alkylsulphinyl, amino $C_{(1-3)}$ alkyl, mono- or di- $C_{(1-3)}$ alkylamino $C_{(1-3)}$ alkyl,  $C_{(1-3)}$ alkylcarbonylamino $C_{(1-3)}$ alkyl,  $C_{(1-3)}$ alkoxy $C_{(1-3)}$ alkylcarbonylamino $C_{(1-3)}$ alkyl,  $C_{(1-3)}$ alkylsulphonylamino $C_{(1-3)}$ alkyl,  $C_{(1-3)}$ alkylcarboxy, or  $C_{(1-3)}$ alkylcarboxy $C_{(1-3)}$ alkyl;

$R^b$  is hydrogen, halogen,  $C_{(1-3)}$ alkyl, or hydroxy $C_{(1-3)}$ alkyl, with the proviso that  $R^a$  and  $R^b$  are not simultaneously each hydrogen; or

$R^a$  and  $R^b$  together are  $(CH_2)_n$  where  $n$  is 3 or 4, to form, with the pyrimidine ring carbon atoms to which they are attached a fused 5- or 6-membered carbocyclic ring; or

$R^a$  and  $R^b$  together with the pyrimidine ring carbon atoms to which they are attached form a fused benzo or heteroaryl ring optionally substituted by 1, 2, 3 or 4 substituents which may be the same or different selected from halogen,  $C_{(1-4)}$ alkyl, cyano,  $C_{(1-4)}$ alkoxy or  $C_{(1-4)}$ alkylthio, or mono to perfluoro- $C_{(1-4)}$ alkyl);

$R^c$  is hydrogen or  $C_{(1-3)}$ alkyl;

$R^2$  is an aryl or heteroaryl group, optionally substituted by 1, 2, 3 or 4 substituents which may be the same or different selected from  $C_{(1-18)}$ alkyl,  $C_{(1-18)}$ alkoxy,  $C_{(1-18)}$ alkylthio, aryl $C_{(1-18)}$ alkoxy, hydroxy, halogen, CN,  $COR^6$ , carboxy,  $COOR^6$ ,  $NR^6COR^7$ ,  $CONR^8R^9$ ,  $SO_2NR^8R^9$ ,  $NR^6SO_2R^7$ ,  $NR^8R^9$ , mono to perfluoro- $C_{(1-4)}$ alkyl, mono to perfluoro- $C_{(1-4)}$ alkoxyaryl, and aryl $C_{(1-4)}$ alkyl;

$R^3$  is hydrogen,  $C_{(1-6)}$ alkyl which may be unsubstituted or substituted by 1, 2 or 3 substituents selected from hydroxy, halogen,  $OR^6$ ,  $COR^6$ , carboxy,  $COOR^6$ ,  $CONR^8R^9$ ,  $NR^8R^9$ ,  $NR^8COR^9$ , mono- or di-(hydroxy $C_{(1-6)}$ alkyl)amino and N-hydroxy $C_{(1-6)}$ alkyl-N- $C_{(1-6)}$ alkylamino; or

$R^3$  is Het- $C_{(0-4)}$ alkyl in which Het is a 5- to 7- membered heterocyclyl ring comprising N and optionally O or S, bonded through a carbon ring atom and in which N may be substituted by  $COR^6$ ,  $COOR^6$ ,  $CONR^8R^9$ , or  $C_{(1-6)}$ alkyl optionally substituted by 1, 2 or 3 substituents selected from hydroxy, halogen,  $OR^6$ ,  $COR^6$ , carboxy,  $COOR^6$ ,  $CONR^8R^9$  or  $NR^8R^9$ ;

$R^4$  is an aryl or a heteroaryl ring optionally substituted by 1, 2, 3 or 4 substituents which may be the same or different selected from  $C_{(1-18)}$ alkyl,  $C_{(1-18)}$ alkoxy,  $C_{(1-18)}$ alkylthio, aryl $C_{(1-18)}$ alkoxy, hydroxy, halogen, CN,  $COR^6$ , carboxy,  $COOR^6$ ,  $NR^6COR^7$ ,  $CONR^8R^9$ ,  $SO_2NR^8R^9$ ,  $NR^6SO_2R^7$ ,  $NR^8R^9$ , mono to perfluoro- $C_{(1-4)}$ alkyl and mono to perfluoro- $C_{(1-4)}$ alkoxy;

$R^5$  is an aryl or heteroaryl ring which is further optionally substituted by 1, 2, 3 or 4 substituents which may be the same or different selected from  $C_{(1-18)}$ alkyl (preferably  $C_{(1-6)}$ alkyl),  $C_{(1-18)}$ alkoxy (preferably  $C_{(1-6)}$ alkoxy),  $C_{(1-18)}$ alkylthio (preferably  $C_{(1-6)}$ alkylthio), aryl $C_{(1-18)}$ alkoxy (preferably

arylC<sub>(1-6)</sub>alkoxy), hydroxy, halogen, CN, COR<sup>6</sup>, carboxy, COOR<sup>6</sup>, CONR<sup>8</sup>R<sup>9</sup>, NR<sup>6</sup>COR<sup>7</sup>, SO<sub>2</sub>NR<sup>8</sup>R<sup>9</sup>, NR<sup>6</sup>SO<sub>2</sub>R<sup>7</sup>, NR<sup>8</sup>R<sup>9</sup>, mono to perfluoro-C<sub>(1-4)</sub>alkyl and mono to perfluoro-C<sub>(1-4)</sub>alkoxy;

R<sup>6</sup> and R<sup>7</sup> are independently hydrogen or C<sub>(1-20)</sub>alkyl, for instance C<sub>(1-4)</sub>alkyl;

R<sup>8</sup> and R<sup>9</sup> which may be the same or different is each selected from hydrogen, C<sub>(1-12)</sub>alkyl; or

R<sup>8</sup> and R<sup>9</sup> together with the nitrogen to which they are attached form a 5- to 7 membered ring optionally containing one or more further heteroatoms selected from oxygen, nitrogen and sulphur, and optionally substituted by one or two substituents selected from hydroxy, oxo, C<sub>(1-4)</sub>alkyl, C<sub>(1-4)</sub>alkylCO, aryl, or aralkyl; or

R<sup>8</sup> and R<sup>9</sup> which may be the same or different is each selected from CH<sub>2</sub>R<sup>10</sup>, CHR<sup>11</sup>CO<sub>2</sub>H or a salt thereof in which:

R<sup>10</sup> is COOH or a salt thereof, COOR<sup>12</sup>, CONR<sup>6</sup>R<sup>7</sup>, CN, CH<sub>2</sub>OH or CH<sub>2</sub>OR<sup>6</sup>;

R<sup>11</sup> is an amino acid side chain such as CH<sub>2</sub>OH from serine;

R<sup>12</sup> is C<sub>(1-4)</sub>alkyl or a pharmaceutically acceptable *in vivo* hydrolysable ester group;

n is an integer from 1 to 4;

X is O or S;

Y is (CH<sub>2</sub>)<sub>p</sub>(O)<sub>q</sub> in which p is 1, 2 or 3 and q is 0 or p is 2 or 3 and q is 1; and

Z is O or a bond.

2. A compound as claimed in claim 1 in which R<sup>a</sup> include chloro, bromo, methyl, ethyl, n-propyl, methoxy, hydroxymethyl, hydroxyethyl, methylthio, methylsulphinyl, aminoethyl, dimethylaminomethyl, acetylaminoethyl, 2-(methoxyacetamido)ethyl, mesylaminoethyl, ethylcarboxy, methanesulfonamidoethyl, (methoxyacetamido)ethyl and iso-propylcarboxymethyl.

3. A compound as claimed in claim 1 or 2 in which R<sup>b</sup> is hydrogen or methyl.

4. A compound as claimed in claim 1 in which R<sup>a</sup> and R<sup>b</sup> together with the pyrimidine ring carbon atoms to which they are attached form a fused 5-membered carbocyclic ring or a fused benzo or heteroaryl ring ring selected from benzo, pyrido and thieno, respectively.

5. A compound as claimed in any one of claims 1 to 4 in which R<sup>c</sup> is hydrogen or methyl.

6. A compound as claimed in any one of claims 1 to 5 in which X is S.


7. A compound as claimed in any one of claims 1 to 6 in which Y is CH<sub>2</sub>.

8. A compound as claimed in any one of claims 1 to 7 in which Z is a direct bond.

9. A compound as claimed in any one of claims 1 to 8 in which R<sup>2</sup> is an aryl group, optionally substituted by 1, 2, 3 or 4 substituents which may be the same or different selected from C<sub>(1-6)</sub>alkyl, C<sub>(1-6)</sub>alkoxy, C<sub>(1-6)</sub>alkylthio, hydroxy, halogen, CN, mono to perfluoro-C<sub>(1-4)</sub>alkyl, mono to perfluoro-C<sub>(1-4)</sub>alkoxyaryl, and arylC<sub>(1-4)</sub>alkyl.

10. A compound as claimed in any one of claims 1 to 9 in which R<sup>2</sup> is phenyl optionally substituted by halogen.

11. A compound as claimed in any one of claims 1 to 10 in which  $R^2$  is phenyl optionally substituted by one to three fluorine atoms.
12. A compound as claimed in any one of claims 1 to 11 in which  $R^3$  is  $C_{(1-3)}$ alkyl substituted by a substituent selected from  $NR^8R^9$ ; or  $R^3$  is Het- $C_{(0-2)}$ alkyl in which Het is a 5- to 7- membered heterocyclyl ring comprising N and in which N may be substituted by  $C_{(1-6)}$ alkyl.
13. A compound as claimed in any one of claims 1 to 12 in which  $R^3$  is 2-(diethylamino)ethyl.
14. A compound as claimed in any one of claims 1 to 13 in which  $R^4$  is phenyl.
15. A compound as claimed in any one of claims 1 to 14 in which  $R^5$  is phenyl substituted by trifluoromethyl, preferably at the 4-position.
16. A compound as claimed in any one of claims 1 to 15 in which  $R^4$  and  $R^5$  together form a 4-(4-trifluoromethylphenyl)phenyl moiety.
17. A compound of the formula (IA):




(IA)

in which:

$R^a$ ,  $R^b$ ,  $R^c$ ,  $n$ ,  $R^2$ ,  $R^3$ ,  $R^4$ ,  $R^5$ , and  $X$  are as defined in claim 1.

18. A compound of the formula (IB):



(IB)

in which:

$R^a$  and  $R^b$  together with the pyrimidine ring carbon atoms to which they are attached form a fused 5-membered carbocyclic ring;

$R^2CH_2X$  is 4-fluorobenzylthio;

$R^3$  is  $C_{(1-3)}$ alkyl substituted by  $NR^8R^9$ ; or

$R^3$  is  $Het-C_{(0-2)}$ alkyl in which Het is a 5- to 7- membered heterocyclyl ring comprising N and in which N may be substituted by  $C_{(1-6)}$ alkyl;

5  $R^4$  and  $R^5$  form a 4-(4-trifluoromethylphenyl)phenyl moiety;

$R^8$  and  $R^9$  which may be the same or different is each selected from hydrogen, or  $C_{(1-6)}$ alkyl; and X is S.

19. A compound as claimed in claim 1 selected from:

- 10 1-(N-(2-(Diethylamino)ethyl)-N-(4-(4-trifluoromethylphenyl)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-ethylpyrimidin-4-one;
- 1-(N-(2-(Diethylamino)ethyl)-N-(2-(4-trifluoromethylphenyl)pyrid-5-ylmethyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-ethylpyrimidin-4-one; 1-(N-(2-(Diethylamino)ethyl)-N-(4-(4-trifluoromethylphenyl)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5,6-trimethylenepyrimidin-4-one;
- 15 1-(N-(2-(Diethylamino)ethyl)-N-(2-(4-trifluoromethylphenyl)pyrid-5-ylmethyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5,6-trimethylenepyrimidin-4-one;
- 1-(N-(2-(Diethylamino)ethyl)-N-(2-(4-trifluoromethylphenyl)pyrimid-5-ylmethyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5,6-trimethylenepyrimidin-4-one;
- 20 1-(N-Methyl-N-(2-(4-chlorophenyl)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(2-aminoethyl)pyrimidin-4-one;
- 1-(N-Methyl-N-(2-(4-chlorophenyl)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(2-acetamidoethyl)pyrimidin-4-one;
- 1-(N-(2-(Diethylamino)ethyl)-N-(2-(4-chlorophenyl)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(dimethylaminomethyl)pyrimidin-4-one;
- 25 1-(N-(2-(Diethylamino)ethyl)-N-(4-(4-chlorophenyl)benzyl)-aminocarbonyl-methyl)-2-(4-fluorobenzyl)thio-5-methylpyrimidin-4-one;
- 1-(N-methyl-N-(2-(4-trifluoromethylphenyl)pyrid-5-ylmethyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-methylpyrimidin-4-one;
- 1-(N-(2-(Diethylamino)ethyl)-N-(4-(4-trifluoromethylphenyl)benzyl)-aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-methylpyrimidin-4-one;
- 30 1-(N-(2-(Diethylamino)ethyl)-N-(2-(4-trifluoromethylphenyl)pyrid-5-yl-methyl)-aminocarbonyl-methyl)-2-(4-fluorobenzyl)thio-5-methylpyrimidin-4-one;
- 1-(N-(2-(Diethylamino)ethyl)-N-(2-(4-chlorophenyl)pyrimid-5-yl-methyl)-aminocarbonyl-methyl)-2-(4-fluorobenzyl)thio-5-methylpyrimidin-4-one;
- 35 1-(N-(2-(Diethylamino)ethyl)-N-(2-(4-trifluoromethylphenyl)pyrimid-5-yl-methyl)aminocarbonyl-methyl)-2-(4-fluorobenzyl)thio-5-methylpyrimidin-4-one;
- (±)-1-(N-(2-(Diethylamino)ethyl)-N-(1-(4-(4-chlorophenyl)phenyl)ethyl)-aminocarbonyl-methyl)-2-(4-fluorobenzyl)thio-5-methylpyrimidin-4-one;
- 1-(N-(2-(1-piperidino)ethyl)-N-(4-(4-trifluoromethylphenyl)benzyl)-aminocarbonyl-methyl)-2-(4-fluorobenzyl)thio-5-methylpyrimidin-4-one;
- 40 1-(N-methyl-N-(4-(4-trifluoromethylphenyl)benzyl)-aminocarbonyl-methyl)-2-(4-fluorobenzyl)thio-5-ethylpyrimidin-4-one
- 1-(N-(2-(Diethylamino)ethyl)-N-(4-(4-chlorophenyl)benzyl)-aminocarbonyl-methyl)-2-(4-fluorobenzyl)thio-5-ethylpyrimidin-4-one;
- 45 1-(N-methyl-N-(2-(4-trifluoromethylphenyl)pyrid-5-yl-methyl)-aminocarbonyl-methyl)-2-(4-fluorobenzyl)thio-5-ethylpyrimidin-4-one;

- 1-(N-(2-(Diethylamino)ethyl)-N-(2-(4-chlorophenyl)pyrimid-5-yl-methyl)-aminocarbonyl-methyl)-2-(4-fluorobenzyl)thio-5-ethylpyrimidin-4-one;
- 1-(N-(2-(Diethylamino)ethyl)-N-(2-(4-trifluoromethylphenyl)pyrimid-5-yl-methyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-ethylpyrimidin-4-one;
- 5 1-(N-(2-(Diethylamino)ethyl)-N-(4-(4-chlorophenyl)benzyl)-aminocarbonyl-methyl)-2-(4-fluorobenzyl)thio-5-propylpyrimidin-4-one;
- 1-(N-(2-(Diethylamino)ethyl)-N-(4-(4-trifluoromethylphenyl)benzyl)-aminocarbonyl-methyl)-2-(4-fluorobenzyl)thio-5-propylpyrimidin-4-one;
- 1-(N-(2-(Diethylamino)ethyl)-N-(4-(4-trifluoromethylphenyl)benzyl)-aminocarbonyl-methyl)-2-(4-fluorobenzyl)thio-5-ethoxycarbonylmethylpyrimidin-4-one;
- 10 1-(N-(2-(Diethylamino)ethyl)-N-(4-(4-trifluoromethylphenyl)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-isopropoxycarbonylmethylpyrimidin-4-one;
- 1-(N-(2-(Diethylamino)ethyl)-N-(4-(4-chlorophenyl)benzyl)-aminocarbonyl-methyl)-2-(4-fluorobenzyl)thio-5-hydroxymethylpyrimidin-4-one;
- 15 1-(N-(2-(Diethylamino)ethyl)-N-(4-(4-trifluoromethylphenyl)benzyl)-aminocarbonyl-methyl)-2-(4-fluorobenzyl)thio-5-hydroxymethylpyrimidin-4-one;
- 1-(N-methyl-N-(4-(4-chlorophenyl)benzyl)-aminocarbonyl-methyl)-2-(4-fluorobenzyl)thio-5-(2-hydroxyethyl)pyrimidin-4-one;
- 1-(N-(2-(Diethylamino)ethyl)-N-(4-(4-chlorophenyl)benzyl)-aminocarbonyl-methyl)-2-(4-fluorobenzyl)thio-5-(2-hydroxyethyl)pyrimidin-4-one;
- 20 1-(N-(2-(Diethylamino)ethyl)-N-(2-(4-trifluoromethylphenyl)pyrid-5-yl-methyl)-aminocarbonyl-methyl)-2-(4-fluorobenzyl)thio-5-(2-hydroxyethyl)pyrimidin-4-one;
- 1-(N-(2-(Diethylamino)ethyl)-N-(4-(4-trifluoromethylphenyl)benzyl)aminocarbonyl-methyl)-2-(4-fluorobenzyl)thio-5-(2-hydroxyethyl)pyrimidin-4-one;
- 25 1-(N-(2-(Diethylamino)ethyl)-N-(4-(4-trifluoromethylphenyl)benzyl)aminocarbonyl-methyl)-2-(4-fluorobenzyl)thio-5,6-dimethylpyrimidin-4-one;
- 1-(N-(2-(Diethylamino)ethyl)-N-(4-(4-chlorophenyl)benzyl)-aminocarbonyl-methyl)-2-(4-fluorobenzyl)thio-5,6-trimethylenepyrimidin-4-one;
- 1-(N-(2-(Diethylamino)ethyl)-N-(4-(4-chlorophenyl)benzyl)-aminocarbonyl-methyl)-2-(4-fluorobenzyl)thio-5,6-tetramethylenepyrimidin-4-one;
- 30 1-(N-(2-(Diethylamino)ethyl)-N-(4-(4-trifluoromethylphenyl)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5,6-tetramethylenepyrimidin-4-one;
- 1-(N-(2-(Diethylamino)ethyl)-N-(2-(4-trifluoromethylphenyl)pyrid-5-yl-methyl)-aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5,6-tetramethylenepyrimidin-4-one;
- 35 1-(N-(2-(Diethylamino)ethyl)-N-(4-(4-trifluoromethylphenyl)benzyl)aminocarbonyl-methyl)-2-(4-fluorobenzyl)thio-5-chloropyrimidin-4-one;
- 1-(N-(2-(Diethylamino)ethyl)-N-(4-(4-chlorophenyl)benzyl)-aminocarbonyl-methyl)-2-(4-fluorobenzyl)thio-5-chloropyrimidin-4-one;
- 1-(N-(2-(Diethylamino)ethyl)-N-(2-(4-trifluoromethylphenyl)pyrid-5-yl-methyl)-aminocarbonyl-methyl)-2-(4-fluorobenzyl)thio-5-chloropyrimidin-4-one;
- 40 1-(N-(2-(Diethylamino)ethyl)-N-(4-(4-trifluoromethylphenyl)benzyl)aminocarbonyl-methyl)-2-(4-fluorobenzyl)thio-5-bromopyrimidin-4-one;
- 1-(N-(2-(Diethylamino)ethyl)-N-(4-(4-chlorophenyl)benzyl)-aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-bromopyrimidin-4-one;
- 45 1-(N-(2-(Diethylamino)ethyl)-N-(4-(4-trifluoromethylphenyl)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-methoxypyrimidin-4-one;

- 1-(N-(2-(Diethylamino)ethyl)-N-(2-(4-trifluoromethylphenyl)pyrid-5-yl-methyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-methoxypyrimidin-4-one;
- 1-(N-(2-(Diethylamino)ethyl)-N-(4-(4-trifluoromethylphenyl)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-ethoxypyrimidin-4-one;
- 5 1-(N-(2-(Diethylamino)ethyl)-N-(2-(4-trifluoromethylphenyl)pyrid-5-yl-methyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-ethoxypyrimidin-4-one;
- 1-(N-(2-(Diethylamino)ethyl)-N-(2-(4-trifluoromethylphenyl)pyrid-5-yl-methyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-methylthiopyrimidin-4-one;
- 1-(N-(2-(Diethylamino)ethyl)-N-(4-(4-trifluoromethylphenyl)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-methylthiopyrimidin-4-one;
- 10 1-(N-(2-(Diethylamino)ethyl)-N-(4-(4-trifluoromethylphenyl)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-methylsulfinylpyrimidin-4-one;
- 1-(N-(2-(Diethylamino)ethyl)-N-(2-(4-trifluoromethylphenyl)pyrid-5-yl-methyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-methylsulfinylpyrimidin-4-one;
- 15 1-(N-(2-(Diethylamino)ethyl)-N-(4-(4-trifluoromethylphenyl)benzyl)amino-carbonylmethyl)-2-(2,3-difluorobenzyl)-thio-5,6-trimethylenepyrimidin-4-one;
- 1-(N-(2-(Diethylamino)ethyl)-N-(4-(4-trifluoromethylphenyl)benzyl)amino-carbonylmethyl)-2-(3,4-difluorobenzyl)-thio-5,6-trimethylenepyrimidin-4-one;
- 1-(N-(2-(Diethylamino)ethyl)-N-(4-(4-trifluoromethylphenyl)benzyl)amino-carbonylmethyl)-2-(2,3,4-trifluorobenzyl)-thio-5,6-trimethylenepyrimidin-4-one;
- 20 1-(N-(2-(Diethylamino)ethyl)-N-(4-(4-trifluoromethylphenyl)benzyl)aminocarbonylmethyl)-2-(2-fluorobenzyl)thio-5,6-trimethylenepyrimidin-4-one;
- 1-(N-methyl-N-(4-(4-trifluoromethylphenyl)benzyl)-aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5,6-trimethylenepyrimidin-4-one;
- 25 1-(N-(2-(1-piperidino)ethyl)-N-(4-(4-trifluoromethylphenyl)benzyl)-aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5,6-trimethylenepyrimidin-4-one;
- 1-(N-(2-(Diethylamino)ethyl)-N-(3-(4-trifluoromethylphenoxy)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5,6-trimethylenepyrimidin-4-one;
- 1-(N-(2-(Diethylamino)ethyl)-N-(4-(4-trifluoromethylphenoxy)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5,6-trimethylenepyrimidin-4-one;
- 30 1-(N-(2-(Diethylamino)ethyl)-N-(4-(4-trifluoromethyl-biphenyl-4-yl)propyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5,6-trimethylenepyrimidin-4-one;
- 1-(N-(2-(Diethylamino)ethyl)-N-(4-(4-trifluoromethyl-biphenyl-4-yl)propyl)-aminocarbonyl-methyl)-2-(4-fluorobenzyl)thio-5-ethylpyrimidin-4-one;
- 35 1-(N-(2-(Diethylamino)ethyl)-N-(4-(4-trifluoromethyl-biphenyl-4-yloxy)ethyl)-aminocarbonyl-methyl)-2-(4-fluorobenzyl)thio-5-ethylpyrimidin-4-one;
- 1-(N-(1-Ethyl-piperidin-4-yl)-N-(4-(4-trifluoromethylphenyl)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5,6-trimethylenepyrimidin-4-one;
- 1-(N-(2-Ethylamino-2-methylpropyl)-N-(4-(4-trifluoromethylphenyl)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5,6-trimethylenepyrimidin-4-one;
- 40 N-(2-tert-butylaminoethyl)-N-(4-(4-trifluoromethylphenyl)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5,6-trimethylenepyrimidin-4-one;
- N-(2-Diethylaminoethyl)-2-[2-(4-fluorobenzylthio)-4-oxo-4 *H*-thieno[3,2-*d*]pyrimidin-1-yl]-N-(4'-trifluoromethyl-biphenyl-4-ylmethyl)-acetamide;
- 45 N-(2-Diethylaminoethyl)-2-[2-(4-fluorobenzylthio)-4-oxo-4 *H*-quinazolin-1-yl]-N-(4'-trifluoromethyl-biphenyl-4-ylmethyl)-acetamide;



Ethyl- {2-[ {2-(4-fluorobenzylthio)-4-oxo-4,5,6,7-tetrahydrocyclopentapyrimidin-1-yl]-ethanoyl} - {4'-trifluoromethylbiphenyl-4-ylmethyl)-amino]-ethyl} carbamic acid tert-butyl ester;

1-(N-(1-Methylpiperidin-4-yl)-N-(4-(4-trifluoromethylphenyl)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5,6-trimethylenepyrimidin-4-one;

5 1-(N-(1-Isopropylpiperidin-4-yl)-N-(4-(4-trifluoromethylphenyl)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5,6-trimethylenepyrimidin-4-one;

1-(N-(1-(2-Methoxyethyl)piperidin-4-yl)-N-(4-(4-trifluoromethylphenyl)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5,6-trimethylenepyrimidin-4-one;

10 1-(N-(2-(Diethylamino)ethyl)-N-(4-(4-trifluoromethylphenyl)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(2-aminoethyl)pyrimidin-4-one; 1-(N-(2-(Diethylamino)ethyl)-N-(4-(4-trifluoromethylphenyl)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(2-acetamidoethyl)-pyrimidin-4-one;

1-(N-(2-(Diethylamino)ethyl)-N-(4-(4-trifluoromethylphenyl)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(2-methanesulfonamidoethyl)pyrimidin-4-one;

15 1-(N-(2-(Diethylamino)ethyl)-N-(4-(4-trifluoromethylphenyl)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5-(2-(methoxyacetamido)ethyl)pyrimidin-4-one; and

1-(N-(2-(Ethylamino)ethyl)-N-(4-(4-trifluoromethylphenyl)benzyl)aminocarbonylmethyl)-2-(4-fluorobenzyl)thio-5,6-trimethylenepyrimidin-4-one; or  
a pharmaceutically acceptable salt thereof.

20. A pharmaceutical composition comprising a compound of formula (I) as claimed in claim 1 and a pharmaceutically acceptable carrier.

21. A compound of formula (I) as claimed in claim 1 for use in therapy.


22. The use of a compound of formula (I) as claimed in claim 1 for the manufacture of a medicament for the primary and secondary prevention of acute coronary events.

23. The use as claimed in claim 22 wherein the coronary event is caused by atherosclerosis.

24. The use of a compound of formula (I) as claimed in claim 1 for the manufacture of a medicament for treating a disease state associated with activity of the enzyme Lp-PLA<sub>2</sub>.

25. A process for preparing a compound of formula (I) as defined in claim 1 which process comprises:

(a) reacting a compound of formula (II):



(II)


in which X, n, R<sup>a</sup>, R<sup>b</sup> and R<sup>2</sup> are as hereinbefore defined,  
with a compound of formula (III):



(III)


in which  $R^c$ ,  $R^3$ ,  $R^4$ ,  $R^5$ , Y and Z are as hereinbefore defined; under amide forming conditions;

(b) reacting a compound of formula (IV):



(IV)


10 in which X,  $R^a$ ,  $R^b$  and  $R^2$  are as hereinbefore defined, with a compound of formula (V):



(V)

15 in which n,  $R^3$ ,  $R^4$ ,  $R^5$ ,  $R^c$ , Y and Z are as hereinbefore defined, and  $L^1$  is a leaving group such as halogen, for instance bromo iodo, or triflate in the presence of a base such as a secondary or tertiary amine, for instance di-isopropylethylamine, in an inert solvent such as dichloromethane;

20 (c) when X is S, reacting a compound of formula (VI):



(VI)


25 in which n,  $R^a$ ,  $R^b$ ,  $R^c$ ,  $R^3$ ,  $R^4$ ,  $R^5$ , Y and Z are as hereinbefore defined, with a compound of formula (VII):



(VII)

30 in which  $R^2$  and  $L^1$  are as hereinbefore defined, in the presence of a base such as a secondary or tertiary amine, for instance di-isopropylethylamine, in an inert solvent such as dichloromethane;

(d) when X is O, reacting a compound of formula (VIII):



(VIII)

- 5 in which n, R<sup>a</sup>, R<sup>b</sup>, R<sup>c</sup>, R<sup>3</sup>, R<sup>4</sup>, R<sup>5</sup>, Y and Z are as hereinbefore defined, and L<sup>2</sup> is a leaving group such as halogen or alkylthio, for instance methylthio, with a compound of formula (IX):



10 (IX)

in which R<sup>2</sup> is as hereinbefore defined,

in the presence of a base such as 4-dimethylaminopyridine, in an inert solvent such as pyridine; or

- 15 (e) converting of compound of formula (I) to another compound of formula (I), by functional group modification, using methods well known to those skilled in the art, for example converting a compound of formula (I) in which R<sup>a</sup> is aminoalkyl to a compound of formula (I) in which R<sup>a</sup> is alkylcarbonylaminoalkyl, by reaction with an acylating agent, such as, for example, acetic anhydride.

## INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 01/01515

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D239/56 C07D401/12 C07D239/70 C07D239/26 C07D213/06  
C07D213/54 C07C15/14 A61K31/505 A61P3/06

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D C07C A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

CHEM ABS Data, EPO-Internal, PAJ, WPI Data

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A, P	<p>DATABASE CHEMABS 'Online! CHEMICAL ABSTRACTS SERVICE, COLUMBUS, OHIO, US; BOYD, HELEN F. ET AL: "N-1-Substituted pyrimidin-4-ones: novel, orally active inhibitors of lipoprotein-associated phospholipase A2" retrieved from STN Database accession no. 134:115922 XP004224263 abstract &amp; BIOORG. MED. CHEM. LETT. (2000), 10(22), 2557-2561 ,</p> <p style="text-align: center;">--- -/-</p>	1-25

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

## \* Special categories of cited documents:

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

- \*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- \*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- \*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- \*G\* document member of the same patent family

Date of the actual completion of the international search

1 June 2001

Date of mailing of the international search report

12/06/2001

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Menegaki, F

## INTERNATIONAL SEARCH REPORT

Additional Application No

PCT/EP 01/01515

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 99 24420 A (SMITH STEPHEN ALLAN ; IFE ROBERT JOHN (GB); PINTO IVAN LEO (GB); HI) 20 May 1999 (1999-05-20) cited in the application claims; examples ---	1-25
P,A	WO 00 66567 A (FENWICK ASHLEY EDWARD ; SMITH STEPHEN ALLAN (GB); IFE ROBERT JOHN) 9 November 2000 (2000-11-09) cited in the application claims; examples -----	1-25

## INTERNATIONAL SEARCH REPORT

information on patent family members

International Application No

PCT/EP 01/01515

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9924420 A	20-05-1999	AU 1157599 A EP 1028955 A	31-05-1999 23-08-2000
WO 0066567 A	09-11-2000	AU 4120300 A	17-11-2000